

THE AMERICAN JOURNAL OF PHARMACY

VOL. 106

DECEMBER, 1934

No. 12

EDITORIAL

REGULATING THE SALE OF MEDICINES

CHARLES D. HOWARD, of the New Hampshire Board of Health, preaches from the pulpit of a little periodical called "Health", issued by the department which he so well serves.

For long years he has been the pert exponent of all that is decent and wholesome in the service of public health.

With his pen and his practice he has persistently fought the forces of fraud and corruption in the drug and cosmetic fields.

And he is not just a crusader in words alone, for he deals in deeds as well. The Laboratories of the New Hampshire Board of Health have been his authority and he is girt with their findings whenever he stages a fight.

In the October, 1934, number of his little bulletin he has printed an article dealing with the evils of indiscriminate drug vending and he sees in the spread of this practice a definite danger to public health. For decades, various agencies, such as Dr. Howard's department and the several bureaus of the American Medical Association have conducted campaigns of education designed though not always destined to reach the public. Public school instruction likewise has largely entered into health matters, quackery exposures, etc. State and Federal laws have been applied to curb at least the really rank and dangerous medical frauds, but hosts of these quackeries still persist, and they will continue to exist so long as our laws are as inadequate as they are claimed to be by those best qualified to know.

The tricks of advertising and the tenseness of the times seem to have offset most of the good accomplished by the agencies noted above. Patent medicines are flourishing as they have never flourished before.

And the "indiscriminate vending of drugs" by no means stops with patent medicines. Even the hypnotic and analgesic drugs of

the barbituric class, once considered as strictly prescription materials, are now made as available as air for those who want them.

Department stores and drugless drug stores, and sometimes even the old line pharmacy freely sell these harmful remedies, like any cut price medicine, to all who come to buy them.

The jaded, nervous woman, whose short nights are long and whose long days are short soon finds that "Sleep, the soother of hearts"—can be bought like ribbon at the counter—only that this is as "poppied sleep"—hired, hurt and haunted.

The worried business man, in the red all day, sleeps in the blue with barbital, yet finds that this death of his each day's life sinks his depression still deeper.

The whole thing is wrong!! Valuable drugs these are, when properly administered. In an intelligent physician's hand they are a blessing and boon, but they are much too dangerous for uncontrolled and indiscriminate practice.

According to an article appearing in the Tarrant Co. Medical Society's Bulletin, even the doctor is awakening to the fact that he is used to foster the counter sales of the so-called ethical remedials. We quote from that article:

"The individual doctor is swamped with free samples. He in turn passes them out indiscriminately to his patients, very often not even removing the label from the bottle. In other cases the name of the product is printed on tins or on a metal cap to a bottle or in some other way so that he cannot remove the name. The doctor thereby becomes the most important factor in advertising the product to the layman. In just this way such items as Petrolagar, Citrocarbonate, Agarol, Viosterol, Halibut Liver Oil, etc., have become extremely popularized so that the layman calls for the item without prescription, and frequently the drug stores and department stores run special sales with these items prominently displayed on counters. The lay purchaser is not in the dark as to their use because full instructions are usually given on the package."

This is, of course, not a novel procedure, for most of our proprietaries got their life-start this way.

There is no doubt that the spread of the drug-less drug store, with its definitely unfair advantage over the old-line pharmacy has made cheaper and tawdrier medicines more available to the public.

But by far the most powerful allies of the quack have been the irresponsible advertising media that have been latterly available to him—and that are for the moment, at least, free of all control.

Hasten the day when the freedom of the press and the freedom of the air as well, in matters such as these, will have become a freedom, well policed and orderly.

IVOR GRIFFITH.

Effects of Tobacco Smoking

The possible important physiologic effects of tobacco smoking have been investigated from several angles. Recently, Dill, Edwards and Forbes (1) reported on the relation of smoking to blood sugar, blood lactic acid and metabolism. The subjects for their experiments remained at rest in the fasting state for ninety minutes before smoking and for forty-five minutes afterward. Each subject smoked one cigarette, inhaling the smoke during a period of from five to ten minutes. Capillary blood was obtained twice before smoking, one or more times during smoking, and frequently during the subsequent forty-five minutes. Three samples of venous blood were also obtained. The sugar determinations were made by the micro method of Folin and Malmros. In addition to observations made on ten subjects in the fasting state, similar experiments were carried out on two subjects beginning about three hours after a light breakfast. They concluded that smoking one cigarette produced no change in blood sugar, lactic acid or respiratory quotient. In regard to oxygen consumption, however, there appeared to be significant alterations subsequent to smoking. While the oxygen consumption after smoking may remain unchanged in some subjects, in others it may increase as much as 10 or 15 per cent. Although this increase is of small magnitude and of doubtful general importance, it does indicate that subjects for basal metabolic rate testing should not smoke in the morning during which the test is to be made.

1. Dill, D. B.; Edwards, H. T., and Forbes, W. H.: Tobacco Smoking in Relation to Blood Sugar, Blood Lactic Acid and Metabolism, *Am. J. Physiol.* 109:118 (July) 1934.

WHAT ORGANIZED PHARMACY IS DOING ABOUT IT!!

THE Editor of the AMERICAN JOURNAL OF PHARMACY asks a very pertinent question in his November editorial discussing the work of the Committee on Economic Security—"What is organized Pharmacy doing about it?" He points to the fact that a medical advisory committee has come into being because of pressure from the medical profession for participation in the formulation of the principles upon which the proposed social legislation of the New Deal will rest. He infers that the advice of pharmacists has not been asked and probably will not be asked unless organized pharmacy does something about it. He will be glad † to learn that the inference is wrong although by nearly every available precedent his reasoning is entirely correct.

The Committee on Economic Security, made up of the Secretaries of Labor, Treasury, and Agriculture, the Attorney-General and the Federal Emergency Relief Administrator has been busily engaged for some months with a staff of highly competent scientists in the fields of Sociology, Economics and Medicine in drawing up a comprehensive program to give protection to the individual against the vicissitudes and hazards of modern life—unemployment, accident, sickness, invalidity, old age and premature death.

The extent to which advice has been sought from those who may be interested in or affected by a comprehensive program of social insurance has depended upon the sequence in which various phases of the broad problem have been considered and the probable extent of the proposed legislation at the coming session of Congress. It is worthy of note also that the Medical Advisory Committee is composed of individuals representing a variety of points of view. It is not a committee appointed by "organized medicine" with instructions to insist upon a program worked out by one or more associations of physicians. It is exactly what its name implies, an advisory committee picked by the Government from among leading thinkers and doers in American Medicine, regardless of their political or organizational affiliations, but with emphasis upon their record of constructive achievement and broadness of view with respect to the problems to be solved. Neither the Medical Advisory Committee nor those selected or still to be selected to advise on matters relating to dentistry, hospitals, public health or pharmacy, are the result of pressure from

† He is!

anybody or any group. These advisory agencies have been or will be created upon the suggestion of the technical staff of the Committee on Economic Security, which has fully recognized the need for advice and counsel from professional groups.

As early as last October the writer was consulted by the staff of the Committee on Economic Security on the pharmaceutical phases of health insurance. It was pointed out to him that unemployment insurance and old age pensions would in all likelihood be considered first and health insurance would be a later development. However certain pertinent information was desired and it was provided in due course. At the same time assurances were given that if and when any topic affecting pharmacy reaches the point of possible inclusion in a program to be sponsored by the Committee, an advisory committee of pharmacists will be selected and appointed. Such an advisory committee must of necessity include men or women of wide experience who have given some thought and study to the principles of the New Deal and to the systems of medical care in vogue in the United States and foreign lands.

The editor's question therefore takes on a new significance. "What is organized pharmacy going to do about it?" To be asked to give advice and to have a list of names of pharmacists printed on a page of some report is "recognition" of a kind, but what have we to offer in the way of constructive suggestions or a program for fitting the 55,000 or more pharmacies into our changing social structure? Medical and Dental Associations have been at work on the problems of sickness insurance, group practice and other similar developments for some time. They have financed surveys of systems of sickness insurance in foreign countries and have accumulated a body of information that furnishes a basis for advisory work of a high character. What have we done and what have we to offer?

At the meeting of the National Drug Trade Conference, representative of all branches of pharmacy, held in Washington, D. C., December 5, 1934, the following resolution was introduced:

"The Conference favors a broad spirit of cooperation with the medical and dental professions to the end that matters of mutual professional and economic interest may be considered from the standpoint of the welfare of all of the health professions and the public. In the matter of institutional medical care, group methods of practice, health insurance and various proposals now being advocated to revise the existing methods of providing medical care, the Conference is ready and willing

to supply information and suggestions on all questions with which it is actively concerned, and hereby offers full cooperation to the government and to the medical, dental, nursing and other public health professions."

The resolution could not be passed until the words, "In the matter of institutional medical care, group methods of practice, health insurance and various proposals now being advocated to revise the existing methods of providing medical care," were eliminated, on the ground that the inclusion of these words might be construed as an endorsement of the majority report of the Committee on the Costs of Medical Care.

It must be plain to any serious student of pharmacy that if such foolishness represents "organized pharmacy" it cannot *do* anything about anything. Fortunately, we have within our ranks both the brains and the experience to serve the Government, the profession and the public exceedingly well. It is only necessary to brush aside reactionary influences sufficiently to permit the newly awakened spirit of professional progress to assert itself.

Whether "organized pharmacy" is to be represented now or in the future in national developments affecting the public health and welfare depends entirely upon how rapidly and how well it can develop a viewpoint and a technique that is worthy of professional pharmacy in its finest sense. There never has been and never will be any doubt that *pharmacy* has a place in such developments. The best minds in the Government and in other professions have been most prompt in recognizing that fact.

ROBERT P. FISCHER,
President, American Pharmaceutical Association.

ORIGINAL ARTICLES

ROMAN NUMERALS AND THEIR USES IN EARLY MANUSCRIPTS AND BOOKS

By Charles H. LaWall

THE following information is not new, but it will be interesting and perhaps informative, especially to those who have occasion to consult ancient books and manuscripts. We use the so-called Arabic numerals in our daily work in ignorance of or oblivious to, the fact that they really came from India.

In spite of the fact that the great majority of the world's populations could not pass a test in simple arithmetical calculations, including common and decimal fractions, proportion and percentage, the great and decisive progress of the world has been made by mathematicians.

From the time of the ancient Egyptians who oriented their pyramids and temples to the then dominating constellations, through the period of the Babylonians who used astronomy to serve the ends of astrology, (thus making possible a system of fakery which would capture some of the degraded intelligence of the twentieth century) the mathematicians have been the great individuals who have made progress possible in human thought and human development.

Beginning with Thales and Pythagoras, progressing through the periods of Democritus, Archimedes, Hipparchus, Heron, Claudius Ptolemaeus, and the other ancient writers on mathematical subjects, down to Roger Bacon, Leonardo da Vinci, Copernicus, Stevin, Galileo, Tycho-Brahe, Kepler, Descartes, Newton, Leibniz, Euler, Gauss, LaPlace, etc., the great minds of the world which have influenced human thought and human progress have been primarily mathematically inclined. The media through which and with which they have worked have been numerals.

Who invented numerals? How have they been evolved?

The average individual pays so little attention to the external form of the figures, and so much attention to their relative position, that he cannot tell, without looking closely, whether Arabic or Roman numerals are used on the dial of the watch or clock which he consults daily, or indeed whether any numerals are employed at all.

It is stated by historians that numbers are older than letters, and if mankind had been furnished with five fingers and a thumb, the world's system of arithmetic would have been duodecimal instead of decimal, which would have necessitated new integers for 10 and 11, and what we now call 10 would have been 12, which would have been divisible by 2, 3, and 4 (instead of only by 2 and 5), and the history of mathematics and perhaps of civilization itself would have been completely changed.

The Egyptians and the Babylonians both employed arbitrary symbols for numerals. The Egyptians had symbols for 1000 (symbolized by a frog) and from that on up to ten millions (the symbol for one of the largest numbers was the figure of a man with his arms outstretched in admiration). The Egyptian system passed through the hieroglyphic to the hieratic form and from that was carried to the Phoenicians and finally to the Syrians. By the time numerals arrived in ancient Greece in the time of Solon (about 600 B. C.) letters came to be substituted for arbitrary symbols.

This system in which letters were used is usually referred to as the Herodian system, after the mathematician of that name who wrote upon the subject about 200 A. D. As used by the Greeks, in this system I stood for one, II for five, Δ for ten, H for one hundred, X for one thousand, and M for ten thousand.

Monogrammatic combinations of these symbols indicated multiples, as II with Δ for fifty, etc. The Hebrews and Syrians influenced this Grecian practice and a new system was evolved in which the first nine letters of the alphabet stood for the units and other letters for multiples of the units.

When the Romans came into the picture, carrying the culture and usages of previous ages we find them adopting symbols from both the Greeks and the Etruscans. The "C", which is customarily believed to be derived from "centum", (one hundred), is really an evolution of the Etruscan sign adapted for this purpose.

The "M" which every schoolboy believes to be the initial letter of the Roman "mille" (a thousand) is really an evolution of the Greek Φ, which, from careless transcribing and false analogy, was later identified with the Greek μ (Mu), and transformed into the letter M.

That this theory is correct is substantiated by the fact that in many of the printed works of the sixteenth and seventeenth centuries the printers tried to set the Greek Φ with available type. The near-

est they could come to it was by using a letter C, making CIO for the symbol for 1000, and from this we can easily trace the evolution of the D for five hundred from the IO.

The V, which in Roman numerology stands for five is in reality the half of the Etruscan X, which was the symbol for ten, which theory is supported by the fact that there was no letter X in the Roman alphabet. The L, which stands for fifty, was evolved from the half of the symbol for one hundred—C.

While these are more or less speculative views, the fact remains that up to the latter part of the sixteenth century, many printers used the improvised Greek Φ for one thousand, and the separated IO for five hundred.

These facts are indispensable for the correct understanding and interpretation of book dates and chapter numerals in printed books prior to the eighteenth century.

When and how the Arabic numerals, which we now employ for all common purposes except dates, came into use, will make a separate story for presentation at some future time.

We do feel sorry, though, for the Roman schoolboys who had to use this cumbersome system (which lacked a decimal point and a zero) in calculating the areas of the public squares or the interest upon the national debt incurred by paying tribute to the Goths.

Happy New Year for CIOIOCCCCXXXV!

Cancer Cases Reported From Medicinal Arsenic

Cases of skin cancer caused by arsenic-containing medicines taken for other conditions have caused Drs. Clifford C. Franseen and Grantley W. Taylor of Boston, Mass., to issue a warning to physicians to be very cautious in giving arsenic as medicine.

Nine cases definitely due to arsenic and five more cases probably caused by arsenic are reported by them (*American Journal of Cancer, October*). In two of the cases, the patients had been exposed to arsenic in the form of a spray for fruits and vegetables. But arsenic given as medicine for the relief of skin diseases and blood disorders caused the cancerous condition in the majority of the cases. The arsenic-containing medicine had been taken by some of the patients as long as forty years before the cancerous condition appeared.

Arsenic has been a common constituent of quack cancer pastes, Drs. Franseen and Taylor also pointed out. They hold it has no place in the treatment of cancer.

ALPHA-DINITROPHENOL
ITS PURIFICATION, WITH QUALITATIVE AND
QUANTITATIVE DATA

By John C. Bird, Z. Panciera and E. G. E. Shafer

During the course of experimental work in this laboratory upon certain aromatic nitro compounds and their salts, it was found necessary to prepare 2-4 dinitrophenol in as pure a state as possible using simple procedures.

Having regard to the recent interest in the use of this chemical as a metabolic stimulant (J. A. M. A. 101.3.193. 1933), and its possible use in obesity under adequate medical supervision, it occurred to the authors that certain distinctive qualitative tests and quantitative data evolved in connection with this substance might be of some interest particularly to those who might wish to investigate further its medicinal possibilities.

Technical dinitrophenol is available from certain of the larger chemical manufacturing concerns and is thus obtained as a dull brownish crystalline semi-dry paste containing, according to our figures, approximately 8 per cent. of water and possessing a strong characteristic phenolic odor. After several different methods of purification were tried the following was found to give a practically 100 per cent. pure product.

Purification of Crude Dinitrophenol

600 grams of the crude product are stirred with 4000 cc. of water to a homogeneous suspension. To this are added 560 cc. of 20 per cent. caustic soda solution and the whole warmed on the water bath to 70 degrees-80 degrees C. A dark brown solution results to which are added 100 gms. "Norite" carbon. After about five minutes during which the carbon is kept well suspended by stirring the liquid is filtered rapidly whilst still hot, giving a clear amber filtrate. 300 gms. of glacial acetic acid diluted with an equal volume of water and warmed to a similar temperature are now added all at once with vigorous stirring. The liquid clouds slightly but clears almost immediately.

On standing a short time crystallization rapidly sets in and the dinitrophenol is obtained in fine pale yellow leaflets. When the solution is quite cold the crystals are filtered, washed with a little cold water to remove traces of acetic acid, and dried in air. The yield is 505 gms. (about 90 per cent, or 84 per cent. on the crude weight). Hydrochloric acid may be used in place of acetic, but according to

our results the color of the resulting product appears darker although the mother liquid is lighter.

Sodium Dinitrophenate

For the preparation of the sodium salt a similar procedure gives a very desirable product.

1125 gms. of crude dinitrophenol are mixed with 2200 cc. of 10 per cent. caustic soda solution and well stirred to a homogeneous mass. The reaction should be just alkaline to litmus. Water is now added to give a volume of 7500 cc. and the whole is warmed until in solution. The Norite (about 200 gms.) is then added and the liquid after filtering hot is allowed to crystallize. The sodium salt forms bright golden needles. Yield 90 per cent. (or crude weight).

Alpha-dinitrophenol made as described forms pale yellow crystals of m. p. 114 degrees C., very slightly soluble in cold water, more easily on boiling, easily soluble in alcohol, ether chloroform and benzene.

The substance possesses practically no odor and little immediate taste followed later, however, by a slight bitterness.

It should conform to the following qualitative tests:

1. Gently warm a small quantity (.1 gm.) alpha-dinitrophenol in an open crucible over a low flame. The substance melts to a bright yellow liquid and sublimes unchanged.
2. Slowly heat .2 gm. of the substance in an open crucible. The material catches fire and burns quietly evolving carbon. There should be no residue after heating to redness for 15 minutes.

Prepare a .05 per cent. solution of alpha-dinitrophenol in hot distilled water. Allow to cool. The liquid is perfectly clear, transparent and of light greenish yellow color. It is slightly acid to litmus. This solution may be used for the tests.

3. To 5 cc. of above t. s. add 3 drops concentrated H_2SO_4 . The solution is immediately decolorized. (Distinction from picric acid).
4. To 5 cc. add 2 drops NaOH solution 20 per cent. The color deepens to golden yellow.
5. To 5 cc. of t. s. add 5 drops of a 10 per cent. solution of KCN. No change in color is noted. (Picric acid gives an amber color, finally turning red).
6. To 5 cc. test solution add 1 cc. of solution of sodium hydrosulphite 1 per cent. to which has been added 2 cc. of 20 per

cent. ammonia solution. Color becomes amber to red, intensified by warming. (Distinction from p-nitro phenol of m. p. 114 degrees C.)

7. To 5 cc. t. s. add 5 drops of saturated sodium sulphide solution. No change takes place. (Further distinction from picric acid, which gives amber red, disappearing on adding a further 5 drops of sulphide.)
8. To 2 cc. t. s. add an equal volume of .04 per cent. Methylene Blue solution. A precipitate of blue black flocks, having a metallic bronze lustre occurs. The precipitate is soluble in excess of water, or on heating, to a bright green solution.
9. To 5 cc. of t. s. add 5 drops 10 per cent. Barium Chloride solution. There should be no precipitate. (Distinction from beta-dinitrophenol.)

The sodium salt is of course more soluble in water than the free dinitrophenol, giving a deep orange to red liquid. The chemical characteristics otherwise are as described for the phenol.

Quantitative Data

For analytical control purposes the water (as water of crystallization) and sodium content were determined as follows:

Weigh accurately 1 to 2 grams of sample in a suitable container (using a platinum crucible or dish if sodium is to be determined on the same sample. Heat in an air oven at 160 degrees to 170 degrees C. for at least two hours, or to constant weight. Cool in desiccator. Weigh. Calculate loss of weight as per cent. total water.

Typical figures are given in the following tables:

No. 1. Weight of Sample—2.3290 grams.

Temperature	Time of heating	Period Total loss	
		Loss %	%
95 to 105°C.	4 hrs.	1.40	1.40
95 to 105°C.	4 "	.36	1.76
95 to 105°C.	4 "	.50	2.26
130°C.	4 "	4.39	6.65
130°C.	3 "	.41	7.06
130°C.	5 "	.78	7.84
150°C.	4 "	.15	7.99
150°C.	2 "	...	7.99
150°C.	2 "	...	7.99
Total water content			7.99%

No. 2. Weight of sample—1.7894 grams.		
Temperature	Time of heating	Total loss %
165°C.	4 hrs.	8.05
170°C.	1 "	8.05
Total water content		8.05%


From these figures it is evident that it is necessary to heat to 160 degrees-170 degrees C. in order to eliminate all water of crystallization. The literature gives this compound as containing 1 H₂O or 8.04 per cent. with which the above results agree.

Sodium Determination

The following method for the determination of sodium was found to give concordant results. The manipulation is simple, and was devised after several careful attempts at simple ignition had resulted in explosion, and methods of reduction and precipitation had led to more or less complicated procedures.

Method

Weigh 1 to 2 grams of sample in a platinum crucible or dish of at least 25 cc. capacity (the dehydrated material from the total water determination may be used). Moisten the sample thoroughly with about 5 cc. concentrated sulfuric acid, making sure that the entire sample is moistened with the acid. Place in a "chimney heater" and heat in such a way that the entire surface of the crucible is heated gently and uniformly. Gradually raise the temperature until the material is thoroughly charred and there is no more tendency to "creep." Remove the crucible from the "chimney heater" and place directly over a small open flame, being careful to avoid spattering. Finally ignite strongly to a pinkish white residue. Cool. Moisten residue with a few cc. concentrated sulfuric acid. Again evaporate off the acid and ignite. Cool. Weigh as anhydrous sodium sulfate.



Calculate per cent. Na on basis of the original sample.

The following table gives figures secured on one sample of the sodium salt of 2-4 dinitrophenol when using this method:

No. 3.		
Test No.	Wt. of sample	% Na Found
1	1.3369	10.36
2*	1.7894	10.25
3*	1.0963	10.29
4	1.0982	10.26

Average 10.29%

*These samples were dehydrated before the sodium determinations were made. Tests 1 and 4 were made on original material.

The average of 10.29 agrees fairly well with the calculated figure of 10.27 allowing for 1 molecule of water and indicates the accuracy of the method. Many trials were necessary however before a satisfactory method of heating could be found owing to persistent creeping of the sample over the sides of the crucible. The "chimney heater" was found to serve the purpose satisfactorily and is merely an indirect heating device. An old ether can (5 lb. size) with the top removed was used in the determinations reported. A nichrome triangle held about midway between top and bottom of the can by running the triangle ends through holes in the sides of the can served as support for the crucible. The purpose of the device is to secure uniform heat on all surfaces of the crucible and its contents. It effectually prevents the beforementioned creeping during the early heating period and eliminates the "spattering" which invariably takes place under direct heat. As with salts of many aromatic nitro compounds our experience with this sodium salt shows it to be much more sensitive to heat than the parent nitro body. The latter may be dried and ignited in a crucible over an open flame using a little care, but it was not found possible to determine ash in the sodium salt in this way.

On heating, sodium dinitrophenate passes through three distinct changes. From the yellow of the crystalline salt, the color becomes increasingly red as the temperature is raised, until the substance fuses to a deep purple or black mass. If the heating is rapid explosion follows immediately after this fusion. If heated slowly the molten material appears to "boil" for some time after which explosion occurs with even greater violence. As small an amount as 0.1 gm. of the salt spread out on an inverted crucible lid will pass through the changes described, finally exploding in what appears to be a vertical plane leaving well marked lines of carbon deposit across the crucible lid. It would appear safe however to heat the salt in an air oven or device as described.—Contribution from the Research Division, John Wyeth & Brother, Philadelphia, Pa.

PREPARATION OF TETRAIODOPHTHALIC ANHYDRIDE

By G. W. Perkins and G. P. Quimba

DURING the course of a study of the X-ray absorption of certain phthaleins, it became necessary to prepare tetraiodophthalic anhydride. The introduction of four halogens into a benzene nucleus obviously would require some powerful halogen carrier or catalyst. Gnehm (1) chlorinated phthalic anhydride in the presence of antimony pentachloride at a temperature of 200 degrees C. to produce tetrachlorophthalic acid. Juvalta (2) catalysed the halogenation with iodine and sulfur trioxide in the form of fuming sulfuric acid. Pratt and Perkins (3) have prepared tetrachlorophthalic anhydride after the method of Duvalta, and Pratt and Young (4) and Pfeiffer and Flater (5) have applied the same principle to the preparation of tetrabromophthalic acid.

Pratt and Shupp (6) prepared tetraiodophthalic anhydride by modifying the method of Juvalta to make it suitable to prepare on a laboratory scale. They do not report any other methods or concentrations of acid as being tried other than 50 per cent. fuming sulfuric acid. We have found in our studies that a large excess of free sulfur trioxide is necessary to catalyze the iodination. Less than 10 per cent. yield of the tetraiodophthalic anhydride with other lower halogenated forms was obtained using 20 per cent. fuming sulfuric acid. With ordinary concentrated sulfuric acid negligible yields of iodophthalic acids were obtained although as long a time as nine hours heating was used in some runs. However, we did note that if a small amount of red phosphorus was added to ordinary concentrated sulfuric acid the effect of the phosphorus was to catalyze the iodination to about the same extent as 20 per cent. fuming sulfuric acid.

Pratt and Perkins (7) found that one equivalent of phthalic anhydride and two equivalents of iodine when mixed with 50 per cent. fuming sulfuric acid and heated at a low temperature (75 de-

(Contribution from the Organic Chemistry Laboratory of the Philadelphia College of Pharmacy and Science and the Emery Laboratory of Cancer Research.)

grees C.) for as long as six days or when heated at a higher temperature (up to 200 degrees C.) for a short period, gave only traces of tetraiodophthalic anhydride while mixtures of di- and tri-iodoacids or anhydrides were obtained (60 per cent. yield of purified products). They also noted in iodinating the dichlorophthalic anhydrides that when the 3, 6-dichlorophthalic anhydride was iodinated at a comparatively low temperature that tetraiodophthalic anhydride would be formed.

The general method for making the halogen substituted phthalic anhydrides by the method of Duvalta is to dissolve the phthalic anhydride in 50 per cent. fuming sulfuric acid, maintaining the acid at a moderate temperature while introducing the required amount of halogen, then continuing the heating for several hours depending on the halogen derivative being prepared, followed by cooling, pouring on cracked ice, and purification. Pratt and Perkins also found in preparing the di-iodo derivatives that the ratio of the yield of 3, 4 and 3, 6 varieties was changed with change of temperature of the halogenation and that there was no change in the amount of the 4, 5 isomer formed, but that there was a slight increase in tri-iodophthalic anhydride with the higher temperatures. Pratt et al., do not report yields obtained in preparing the tetrachlorophthalic anhydride and report yields of 68 per cent. of the di-iodophthalic anhydrides. They state that they always obtained good yields of tetraiodophthalic anhydride.

We have made a number of preparations of tetraiodophthalic anhydride and find that the strength of the acid is important. In our preparations we used 60 per cent. fuming sulfuric acid (60% free SO_3) while Pratt et al., used 50 per cent. fuming sulfuric acid after the method of Duvalta. Pratt does not give any definite time for heating the reaction mixture, but states it must be heated until the cessation of sulfur dioxide. Practically, this is rather difficult to determine and we have found that good yields could be obtained with definite time and temperature ranges. The following tabulation shows the results obtained in a number of preparations we have carried out using different quantities, temperatures, and mode of addition of iodine.

TABLE 1

Run	Phthalic Anhydride	60% fuming H ₂ SO ₄	Iodine	C ₆ I ₄ (CO) ₂ O yield	Per cent. yield
1.	5 Gm.	40 Cc.	20 Gm.	13.9 Gm.	63.1
2.	10 Gm.	80 Cc.	40 Gm.	34.0 Gm.	77.2
3.	15 Gm.	120 Cc.	60 Gm.	45.0 Gm.	68.1
4.	15 Gm.	120 Cc.	60 Gm.	30.0 Gm.	45.4
5.	15 Gm.	120 Cc.	60 Gm.	55.0 Gm.	83.2
6.	20 Gm.	160 Cc.	80 Gm.	59.0 Gm.	67.0
7.	5 Gm.	40 Cc.	20 Gm.	20.0 Gm.	90.8
8.	25 Gm.	100 Cc.	90 Gm.	94.0 Gm.	85.4
9.	25 Gm.	100 Cc.	90 Gm.	97.0 Gm.	88.1
10.	25 Gm.	100 Cc.	90 Gm.	85.0 Gm.	77.2
11.	25 Gm.	100 Cc.	90 Gm.	83.0 Gm.	75.4
12.	25 Gm.	100 Cc.	90 Gm.	86.0 Gm.	78.1
13.	25 Gm.	100 Cc.	90 Gm.	85.0 Gm.	77.2
14.	25 Gm.	100 Cc.	90 Gm.	85.0 Gm.	77.2
15.	25 Gm.	100 Cc.	90 Gm.	85.0 Gm.	77.2

NOTES

1-3. Oil bath heated to 100 degrees C. and iodine added over a period of 30 minutes. Temperature kept at 100 degrees C. for 5 hours, then heated to 175 degrees C. for 30 minutes and allowed to cool.

4-6. Iodine added at bath temperature of 110 degrees C. Final temperature 175 degrees C. for 30 minutes after heating for 5 hours at 110 degrees C.

7. Iodine introduced at bath temperature of 100 degrees C., oil bath temperature then raised to 135 degrees C. for 5 hours, raising the temperature to 175 degrees C. and immediately cooling.

8. Iodine introduced slowly at a bath temperature of 90 degrees C. Temperature slowly raised to 175 degrees C. over a period of 2 hours, and held there for 15 minutes before cooling.

9. Same as 8. Iodine added at a faster rate.

10. Same as 8. Iodine added in two portions, approximately 60 per cent. first and balance after 30 minutes.

11. Temperature of bath 100 degrees C. during addition of iodine and held there for 3 hours after all iodine was added, then held at 175 degrees C. for 15 minutes.

12-15. Temperature of oil bath held at 70 degrees C. during the addition of iodine and for 2 hours after, then raised to 175 degrees C. and held there for 15 minutes.

The following procedure was then devised based on the findings in the previous table and on other miscellaneous observations. The procedure is somewhat similar to that of Pratt and Shupp (*loc. cit.*) except for the strength of the acid and the definitely prescribed time for each operation. The *average* yield of the seven runs tabulated below the procedure is 85 per cent. based on the phthalic anhydride. For an oil bath we have found that a ten-quart shallow form ice cream can, set on a single burner gas hot-plate was very satisfactory. This will accommodate several preparations at one

time. The product obtained in these preparations was considered pure enough for the preparation of phthaleins.

Tetraiodophthalic Anhydride.—25 gm. of phthalic anhydride are placed in an 800 cc. Kjeldahl flask, followed by 100 cc. of 60 per cent. free SO_3 fuming sulfuric acid. The flask is suspended in an oil bath heated 95 to 105 degrees C., and 90 gm. of finely ground iodine crystals are added to the flask in 2-5 gram portions giving the flask a swirling motion after the addition of each portion of iodine. (A test-tube filled with water may be suspended in the neck of the flask by means of asbestos paper to condense sublimed iodine.) This operation takes about 30 minutes, after which heating is continued at the same temperature for 2 hours more. Finally over a period of thirty minutes the temperature is raised to 175 degrees C. and held there for 15 minutes, and the flask is raised out of the oil bath and allowed to cool to room temperature. The contents of the flask after cooling is then poured into a 2 liter beaker half full of cracked ice. After stirring vigorously and the ice has partially melted, 500 cc. of water are added and sulfur dioxide bubbled through the liquid until most of the iodine is dissolved. The supernatant liquid is decanted off and a liter of warm water added. More sulfur dioxide is passed through the supernatant liquid until it is colorless. A yellow crystalline precipitate settles out. Part of the supernatant liquid is decanted off and the precipitate filtered on a Büchner funnel. The precipitate is washed on the filter with warm water until the washings are free from acid.

The results of several runs made according to the method outlined above is given in Table 2. In runs 2-4 the quantities were varied from those in the rest of the table. In only one run out of a total of twenty-six reported in this paper were we able to obtain a theoretical yield. Pratt on one occasion reports a theoretical yield was obtained and later says that "average yields were obtained."

TABLE 2

Run	Phthalic Anhydride	60% fuming H_2SO_4	Iodine	$\text{C}_6\text{I}_4(\text{CO})_2\text{O}$ yield	Per cent. yield
16.	25 Gm.	100 Cc.	90 Gm.	90.0 Gm.	81.7
17.	20 Gm.	80 Cc.	72 Gm.	79.0 Gm.	91.7
18.	25 Gm.	100 Cc.	90 Gm.	110.0 Gm.	99.9
19.	40 Gm.	160 Cc.	144 Gm.	144.0 Gm.	81.7
20.	25 Gm.	100 Cc.	90 Gm.	95.0 Gm.	86.3
21.	25 Gm.	100 Cc.	90 Gm.	87.0 Gm.	79.0
22.	25 Gm.	100 Cc.	90 Gm.	95.0 Gm.	86.3
23.	25 Gm.	100 Cc.	90 Gm.	88.0 Gm.	80.0
24.	25 Gm.	100 Cc.	90 Gm.	93.0 Gm.	84.3
25.	25 Gm.	100 Cc.	90 Gm.	95.0 Gm.	86.3
26.	25 Gm.	100 Cc.	90 Gm.	91.0 Gm.	82.7

Control analyses were made on all batches of crude product after drying. A persistent impurity not always removed by the sulfur dioxide was free iodine. Sometimes a sample would have a bright yellow color but on control analysis, would have a high percentage of iodine. These samples on heating in an oven at 120 degrees C. over night would lose the adsorbed iodine by sublimation. Pratt and Shupp purified their product further by forming an alkali salt; reprecipitating with acid; heating to form the anhydride, and recrystallizing from nitrobenzene or phenol. The melting point of their product after purification is given as 320-325 degrees C. (cor). Four samples taken from different batches of our crude products melted 323-326; 325-328; 325-327; 325-328 degrees C. Pratt and his co-workers used the lime combustion method for determining the halogen in most of their derivatives. For our control analyses we have used a fusion method using sodium carbonate and sodium peroxide in a nickel crucible. This method was sufficiently accurate for use as a control on the efficiency of our halogenations. We have also used the method of Stepanov and the Thompson-Oakdale method for the iodine content, which in these cases did not seem to be as satisfactory as the fusion method employed.

Method of Analysis: Weigh 0.2000 to 0.3000 gm. of sample and mix with 1.0 gm. of sodium peroxide and 2 grams of sodium carbonate. Both chemicals are reagent grade. Before placing the mixture in a nickel crucible, weigh about 4 gm. of sodium carbonate and pack in the bottom of a nickel crucible (20 cc. size) so that there will be a depression in the sodium carbonate to receive the prepared sample mixture. The prepared sample mixture is then placed in the crucible in the prepared depression. The rest of the crucible is then packed with 13 to 14 gm. of sodium carbonate, which is firmly packed with a glass rod. The crucible is heated over a low flame for 30 minutes after which the flame is increased and the crucible heated at a cherry red heat for 2 hours. The crucible is allowed to cool to room temperature and carefully dissolved in 450 to 500 cc. of distilled water. Occasionally there is a black residue which is insoluble in the water and is filtered off and the filter thoroughly washed. The washings are added to the filtrate and the whole treated with sulfur dioxide until the solution smells strongly of the sulfur dioxide, 10 per cent. sulfuric acid is then added slowly with constant stirring until acid to litmus. The solution is then heated to boiling and 5 per cent. silver nitrate solution added to precipitate the iodide. Some cloudiness may be present due to sulfur dioxide, this may be cleared up by add-

ing a few cc. of concentrated nitric acid and warming. The solution is digested over a low flame until the silver iodide becomes granular, then filtered through a tared Gooch crucible. The precipitate is thoroughly washed, dried and weighed.

The method of fusion as given above is modified from that of Pitzer (8) who lined the nickel crucible with molten sodium carbonate before adding the sample and peroxide. When the fusion is started at a low heat the loss of iodine through vaporization is negligible, so that for rapid control work it was not considered essential to employ an inverted crucible as in the Piria-Shiff method. Further the limestone used in the latter method is sometimes difficult to dissolve after fusion.

The following analyses represent results obtained on different batches of crude product.

TABLE 3

Grams Sample	Weight of AgI	Per cent. Iodine
1. 0.2000 Gm.	0.2792 Gm.	75.3
0.2000 Gm.	0.2810 Gm.	75.87
2. 0.2000 Gm.	0.2886 Gm.	77.91
0.2000 Gm.	0.2878 Gm.	77.7
3. 0.4133 Gm.	0.5976 Gm.	78.07
0.4786 Gm.	0.6885 Gm.	77.6
4. 0.2000 Gm.	0.2956 Gm.	79.81
0.2000 Gm.	0.2952 Gm.	79.20
5. 0.1000 Gm.	0.1478 Gm.	79.81
0.2000 Gm.	0.2952 Gm.	79.20
6. 0.2000 Gm.	0.2866 Gm.	77.38
0.2000 Gm.	0.2892 Gm.	78.08

There is a possibility of loss of iodine at two steps in the above analysis, first, in the fusion of the sample, second, during the neutralization of the sodium carbonate. When the sulfuric acid is added too rapidly some iodine vapors will volatilize even though sulfur dioxide is present. The Thompson-Oakdale method gave uniformly low results with these halogen compounds and it was difficult to drive all of the iodine over into the absorption flask, we believe the introduction of an air stream into the reaction flask would help this deficiency. In a later paper we will compare the fusion method above, with the Carius method which is more accurate, but much more tedious and time consuming.

Summary

1. An improved method for the preparation of tetraiodophthalic anhydride has been outlined with which average yields of 85 per cent. may be obtained.

2. The use of 60 per cent. free sulfur trioxide fuming sulfuric acid is essential to the production of good yields of the tetraiodophthalic anhydride.

3. A method of analysis suitable for rapid control work on tetraiodophthalic anhydride has been outlined.

BIBLIOGRAPHY

- (1) U. S. Patent 322,369, July 14, 1885.
- (2) German Patent 50,177, Fortschr. Teerfärb.-fabrik, 2, 93 (1899).
- (3) Pratt and Perkins, *J. A. C. S.* 40, 204 (1918).
- (4) Pratt and Young, *J. A. C. S.* 40, 204 (1918).
- (5) Pfeiffer and Flater, *Ber.* 55, 413-429 (1922).
- (6) Pratt and Shupp, *J. A. C. S.* 40, 254-255 (1918).
- (7) Pratt and Perkins, *J. A. C. S.* 40, 221-223 (1918).
- (8) Pitzer, L. E., *Chemist-Analyst* 47, 13 (1926).

REPRINTED ARTICLE

THE PREPARATION OF STERILE SOLUTIONS*

By H. Davis

(From the Department of Bacteriology, University College
Hospital Medical School)

WITH the introduction of the British Pharmacopœia, 1932, official methods of sterilization underwent considerable modification, and the pharmacist was made aware of the inadequacy of the methods which were previously in constant practice. The mere boiling of a solution or steaming at atmospheric pressure were laid aside, and autoclaving, Tyndallization and filtration received official recognition. Results derived from the heat treatment of resistant spores in broths and water apparently justified this conclusion, but there appeared to be little evidence on the effect on micro-organisms of the solutions themselves. With this object in view experiments were commenced; solutions of common hypodermic medicaments were prepared, infected with various types of bacteria, subjected to different methods of sterilization and bacteriologically examined. The results were of such an interesting nature that early publication of the preliminary results seemed advisable in the hope that it might stimulate persons interested in the subject to test and possibly to confirm these results. The writer hopes to continue this subject in a series of papers recording a survey of sterilization as applied to pharmaceutical products.

A synopsis of the sterilization methods of different national pharmacopœias is evidence of the divergence of opinion in regard to the means to be adopted for the production of sterile solutions:

The British Pharmacopœia, 1932:—Autoclaving at 115 degrees to 116 degrees C. (10 lb.) for thirty minutes, Tyndallization in final containers at 80 degrees C. for one hour on three successive days, filtration followed by sterility tests, and an emergency method.

The United States Pharmacopœia, X, 1926:—Autoclaving at 115 degrees C. for thirty minutes, fractional sterilization on three consecutive days in boiling water for fifteen minutes or streaming steam for thirty minutes, and sterilization by filtration.

*Reprinted from *Quart. Jour. of Pharmacy and Pharmacology* (London, England).

The Belgian Pharmacopœia, IV, 1930:—Autoclaving at 120 degrees C., boiling for fifteen minutes, Tyndallization, filtration and preparation with aseptic precautions.

The Swiss Pharmacopœia, V, 1933:—Boiling for twenty minutes, boiling with water for twenty to thirty minutes, heating in free-flowing steam for thirty minutes, autoclaving at 110 degrees to 120 degrees C. for fifteen to twenty minutes, Tyndallization at 60 degrees to 65 degrees C. for one hour on three successive days.

The Italian Pharmacopœia, V, 1927:—Streaming steam for one hour, autoclaving at 112 degrees, 120 degrees and 135 degrees C. for thirty minutes, boiling with water for ninety minutes, Tyndallization at 50 degrees to 60 degrees C. for one hour on five or six successive days.

The German Pharmacopœia, VI, 1926:—Autoclaving at 115 degrees C. for fifteen minutes, boiling with water for thirty minutes, steam sterilizing at atmospheric pressure for thirty minutes, Tyndallization at 70 degrees to 80 degrees C. for forty to sixty minutes on at least four consecutive days the solutions being maintained at about 30 degrees C. in the intervals, and filtration.

The Danish Pharmacopœia, VIII, 1933:—Autoclaving at 120 degrees C. for twenty minutes, streaming steam for one hour, heating at 80 degrees C. for two hours and filtration. The products obtained by methods other than autoclaving, before being decribed as sterile, must undergo sterility tests.

The Dutch Pharmacopœia, V, 1926:—Streaming steam for one hour at 100 degrees C., autoclaving at 107 degrees to 110 degrees C. for thirty minutes, with aqueous liquids by boiling for thirty minutes, Tyndallization at 60 degrees to 65 degrees C. on five consecutive days or at 70 degrees to 80 degrees C. on three consecutive days, and preparation with aseptic precautions.

A lack of unanimity in sterilization methods prescribed by these different authorities is therefore evident.* Autoclaving and Tyndallization are accepted by most as efficient procedures, but there is considerable variation in the other methods. As there appeared to be no doubt of the efficacy of autoclaving and as the result of in-

*While this paper was passing through the press Dr. Hampshire kindly supplied me with the following details from the new *Hungarian Pharmacopœia*, IV, 1934:—Steaming at 100 degrees C. for half an hour, Tyndallization at 70 degrees to 80 degrees C. for forty to sixty minutes on three successive days, the temperature being maintained at 25 degrees to 30 degrees C. between the periods of heating. Details of times and temperatures are varied in special instances.

numerable tests applied to autoclaved products, no further bacteriological investigation of this problem was attempted. Boiling and steaming were then considered. The chief disadvantages in regard to boiling are the necessity to adjust to volume after heating and the enforced use of heat-resisting vessels during the process. Exposure to streaming steam obviates both and an investigation of this process appeared to suggest the most promise.

It is a well-known fact that in the preparation of culture media a single steaming can never be completely relied upon. However, there appeared to be little information in regard to the bacteriological examination of hypodermic solutions submitted to this process. It is obvious from the varied nature of the substances that conditions obtaining in these solutions must greatly differ from those in broths and other culture media. The pH of the solution, concentration and osmotic effects and possible bactericidal properties of the solute must of necessity exert some influence on any organisms present. "The rate of disinfection by hot water is greatly increased if the solution is made either acid or alkaline" (Chick, 1910) (1). With this in view experiments were commenced. Determinations of the pH of a number of injections were made colorimetrically and, as was to be expected, a great divergence between the acid and alkaline limits was found. The following table shows some of the results obtained.

TABLE I

pH	Solutions
7 to 9	25 per cent. caffeine-sodium benzoate (pH 8.0), 20 per cent. hexamine (8.2).
6 to 7	0.12 per cent. atropine sulphate (6.8), normal saline (7.0), 5 and 30 per cent. dextrose (7.0 and 6.6), 0.6 per cent. homatropine hydrobromide (6.4).
5 to 6	2 per cent. procaine hydrochloride (5.8), 0.75 per cent. strychnine hydrochloride (5.5), 2.5 per cent. morphine tartrate (6.0).
4 to 5	5 per cent. codeine phosphate (4.7), 3 per cent. pilocarpine nitrate (4.7), 5 per cent. amylocaine hydrochloride (4.9), 2.5 per cent. morphine hydrochloride (4.8), 30 per cent. sodium salicylate (4.4).

A short series of experiments was then made with three of these. The containers used were 25 mil vaccine bottles of neutral glass (Jena). These were sterilized by dry heat and the solutions made with sterile water and infected with suspensions of *Staphylococcus aureus* (twenty-four hour culture) and of *Bacillus mycoides* and *Bacillus subtilis*. These were steamed for five, ten and fifteen

minutes and bacteriologically tested. The staphylococci were killed in five minutes and owing to the ease with which this occurred all future experiments were made with sporing organisms. The results with the other two are formulated in Tables II and III.

One can see the influence of the pH of the codeine phosphate solution on the heat disinfection of *B. mycoides*. Owing to the inadequacy of these short exposures to steam the experiments were repeated, the times being fifteen, thirty and sixty minutes, allowing

TABLE II

Solution	Organism	5 mins.	Steaming 10 mins.	15 mins.
0.12 per cent.	<i>B. mycoides</i>	+	+	+
Atropine sulphate	<i>B. subtilis</i>	+	+	+
5 per cent. Codeine	<i>B. mycoides</i>	—	—	—
phosphate	<i>B. subtilis</i>	+	+	+
Normal saline solution	<i>B. mycoides</i>	+	+	+
" " "	<i>B. subtilis</i>	+	+	+

an additional ten minutes for the contents to attain the temperature of the steam.

TABLE III

Solution	Organism	15 mins.	Steaming 30 mins.	60 mins.
0.12 per cent.	<i>B. subtilis</i>	+	—	—
Atropine sulphate	<i>B. mycoides</i>	+	—	—
5 per cent. Codeine	<i>B. subtilis</i>	+	—	—
phosphate	<i>B. mycoides</i>	—	—	—
25 per cent. Caffeine	<i>B. subtilis</i>	+	—	—
and sodium benzoate	<i>B. mycoides</i>	—	—	—
Normal saline solutions	<i>B. subtilis</i>	+	—	—
	<i>B. mycoides</i>	+	—	—

In every case sterile products were obtained after thirty and sixty minutes' steaming. The inhibitory effect of the solution on the growth of the organism in the broth is discounted by the growth which occurred in the test broths for these solutions steamed for only fifteen minutes. Approximately 30 mls of broth were used for 1 mil of the solution and where precipitation of the medium occurred a test using an agar plate was also done in addition to sub-inoculation of the precipitated broth. All the tests were examined on each day for a total of seven days. Throughout the whole series of tests in this paper if no growth occurred within three days no growth appeared during the remaining four. The tests were carried out in accordance with the directions of the British Pharmacopœia.

Another series of injections was prepared as before and dust, collected from a storeroom in which glassware is unpacked from straw packing material, etc., was used as a further contamination. Examination of the dust-infected material was made only after sixty minutes' steaming.

TABLE IV

Solution	Organism	15 mins.	Steaming 30 mins.	60 mins.
30 per cent. Dextrose	<i>B. subtilis</i>	+	—	—
	<i>B. mycoides</i>	+	—	—
	Dust	—	—	—
5 per cent. Stovaine	<i>B. subtilis</i>	—	—	—
	<i>B. mycoides</i>	—	—	—
	Dust	—	—	—
10 per cent. Soluble barbitone	<i>B. subtilis</i>	+	—	—
	<i>B. mycoides</i>	—	—	—
	Dust	—	—	—
5 per cent. Calcium chloride	<i>B. subtilis</i>	+	—	—
	<i>B. mycoides</i>	+	—	—
	Dust	—	—	—

The sterilization of hexamine by the application of heat has been the subject of much controversy, Martindale (2) states that no decomposition of a 40 per cent. solution has occurred on heating for twenty minutes at 100 degrees C. Other references are much too numerous to mention and it was decided to include hexamine and morphine tartrate in this series. As previous experiments had shown the cultures of *B. mycoides* to be not very resistant this was omitted.

TABLE V

Solution	Organism	15 mins.	Steaming 30 mins.	60 mins.
20 per cent. Hexamine	<i>B. subtilis</i>	+	—	—
	Dust	+	—	—
10 per cent. Soluble penobarbitone	<i>B. subtilis</i>	+	—	—
	Dust	+	—	—
2.5 per cent. Morphine hydrochloride	<i>B. subtilis</i>	+	—	—
	Dust	—	—	—
2.5 per cent. Morphine tartrate	<i>B. subtilis</i>	+	—	—
	Dust	+	—	—

Soluble phenobarbitone was decomposed during the process; white precipitates rapidly formed in the solution steamed for thirty and sixty minutes and in the course of three days appeared in the one heated for fifteen minutes. Obviously solutions of this substance cannot be subjected to steaming for purposes of sterilization.

A culture of *Clostridium sporogenes* was supplied by Mr. H. Berry as a test organism of the resistant sporing anaerobic type, and this was utilized in the next series instead of the dust which was not as stringent a test as previously anticipated.

TABLE VI

Solution	Organism	15 mins.	Steaming 30 mins.	60 mins.
12 per cent. Sodium thiosulphate	<i>B. subtilis</i>	+	—	—
	<i>Cl. sporogenes</i>	+	+	—
0.75 per cent. Strych- nine hydrochloride	<i>B. subtilis</i>	—	—	—
	<i>Cl. sporogenes</i>	—	—	—
5.0 per cent. Peptone	<i>B. subtilis</i>	+	—	—
	<i>Cl. sporogenes</i>	+	—	—
30 per cent. Sodium salicylate	<i>B. subtilis</i>	+	—	—
	<i>Cl. sporogenes</i>	+	—	—

The interesting feature of these results is the comparative ease with which these organisms were killed in the strychnine solution. This is in agreement with the results of Todd and Sillar (3), and there can be little doubt that strychnine hydrochloride possesses some germicidal power. In the case of sodium thiosulphate solution the anaerobe was not killed with thirty minutes' steaming. It was therefore resolved in the final series of tests to adopt sixty minutes as the period for steaming.

All the previous experiments appeared to support those statements in favor of sterilizing hypodermic solutions by exposure to streaming steam for one hour. The British Pharmaceutical Codex, 1934, suggests heating to 100 degrees C. for one hour for codeine phosphate and presumably this would be done in a steamer. Thirty minutes is prescribed for a number of other drugs. A series of experiments was suggested to confirm the previous results and to compare these with those produced by the official Tyndallization process. Solutions were prepared with distilled water in vessels, scrupulously cleaned but not sterilized. These were transferred to 25 mil vaccine bottles and plugged with non-absorbent cotton wool. The infecting organisms for the steaming process were *B. subtilis*, *Cl. sporogenes* and a filtrate prepared from farmyard soil containing straw, stable and poultry manure. A little of this soil was triturated with water and filtered through a coarse filter paper. After boiling a little for seventy-five minutes the liquid was not sterile. As a control a little was mixed with broth and steamed for sixty minutes. On incubation copious growth occurred. Plating and growth in

anaerobic tubes showed the presence in the unheated filtrate of coliform and subtilis-type bacteria and gas-producing anaerobes. Contamination of injections with this material was therefore considered to be a stringent test, the conditions being much worse than any which would be experienced by a person engaged in their preparation. The pH of the solutions, both before and after steaming, was determined as appreciable changes would probably indicate a certain degree of decomposition. The subject of decomposition must be treated as a different problem and this paper is concerned essentially with the *bacteriological* control of solutions.

TABLE VII

Solution	Soil Filtrate		Steamed for 60 mins. <i>B. subtilis</i> & <i>Cl. sporogenes</i>		Tyndallized <i>B. subtilis</i> & <i>Cl. sporogenes</i>		pH	
	Aerobic	Anaerobic	Aerobic	Anaerobic	Aerobic	Anaerobic	Before steaming	After steaming
0.12 per cent. Atropine sulphate	—	—	—	—	+	+	6.6	6.4
5 per cent. Codeine phosphate	—	—	—	—	+	+	4.7	4.7
25 per cent. Caffeine sodium benzoate	—	—	—	—	+	+	8.0	8.0
Normal saline solution	—	—	—	—	+	+	7.0	7.0
5 per cent. Dextrose	—	—	—	—	+	+	7.0	6.0
0.6 per cent. Homatropine HBr.	—	—	—	—	+	+	6.4	6.0
2 per cent. Procaine HCl	—	—	—	—	+	+	6.0	4.2
3 per cent. Pilocarpine nitrate	—	—	—	—	+	—	4.7	4.2

The remarkable fact is evidenced that the Tyndallization process of maintaining at 80 degrees C. for one hour on three successive days is useless when applied to spore infected material, whereas steaming for one hour in every case produced a sterile product. In view of the British Pharmacopœia not specifying the application of sterility tests to Tyndallized products it is evident that in order to place any reliance on the method of solutions *should be prepared with aseptic precautions in previously sterilized containers* before submitting them to the stages of successive heatings. In the remaining series a further step was introduced into the Tyndallization process, a duplicate batch being incubated at 37 degrees C. between the periods intervening between the successive heatings. The temperature inside

the oven was checked by inserting a thermometer through the rubber cap of a vaccine bottle containing water, allowing the temperature of this to reach 80 degrees and maintaining at 80 degrees to 85 degrees C. for one hour. A thermometer in the oven itself was no indication of the temperature attained by the solution.

TABLE VIII

Solution	Steaming for 60 mins.		Tyndallized		<i>B. subtilis</i> & <i>Cl. sporogenes</i>		pH	
	<i>B. subtilis</i> & <i>Cl. sporogenes</i>		Without incubation		With incubation		Before steaming	After steaming
	Aerobic	Anaerobic	Aerobic	Anaerobic	Aerobic	Anaerobic		
20 per cent. Hexamine	—	—	—	—	—	—	8.2	9.4
5 per cent. Amylocaine HCl.	—	—	—	—	—	—	4.9	4.2
10 per cent. Medinal	—	—	—	—	+	—	9.4	9.4
30 per cent. Dextrose	—	—	—	—	+	—	6.6	4.6
5 per cent. Calcium chloride	—	—	—	—	+	—	7.2	7.4
12 per cent. Sodium thiosulphate	—	—	—	—	+	—	7.4	7.4
2.5 per cent. Morphine HCl.	—	—	—	—	+	+	4.8	4.8
2.5 per cent. Morphine tartrate	—	—	—	—	+	+	6.0	6.0
5 per cent. Peptone	—	—	—	—	+	+	6.0	5.8
0.75 per cent. Strychnine HCl.	—	—	—	—	+	+	5.6	5.6
40 per cent. Phenazone	—	—	—	—	—	+	6.4	7.0
30 per cent. Sodium salicylate	—	—	—	—	—	+	4.4	4.4

Because of the negative results of the anaerobic tests on the first six injections in Table VIII a test was made on the broth culture itself. The result showed that the organism had died in this *without* the application of heat, although it was a living culture when introduced into the solutions. Unfortunately it was not possible to repeat these tests with a fresh culture owing to the limited time available for the completion of this paper. They will be repeated and reported in a further communication. In the remainder a fresh culture was used with the results shown. The aerobic tests on the first six demonstrated that only two were sterile, and it is highly probable that when the anaerobic tests are repeated the result will be different

from those previously obtained. The solution of hexamine was sterile after Tyndallization. This supports the views of Pfister (4) that the decomposition products render the solution sterile after a slight degree of heating. That decomposition does occur to some extent on steaming is evidenced by the change in pH from 8.2 to 9.4 and the positive result with Nessler's solution on the steamed product. The test on the unheated control gave a negative result. The results in the case of procaine hydrochloride, pilocarpine nitrate, dextrose and amylocaine hydrochloride appear to indicate some decomposition. The change in pH of procaine hydrochloride confirms the results previously published by Rae, but the writer is not convinced that the degree of decomposition is sufficient to condemn this method of sterilization. For many years solutions of this salt heated to 100 degrees C. have been used by anaesthetists, and in the writer's personal experience the anaesthetic action of these preparations has never been questioned. In regard to amylocaine hydrochloride there appears to be more justification for condemnation. In this case the activity of solutions heated to 100 degrees C. or above has been queried, but since the application of the official Tyndallization process there has been no criticism. The above statement is the result of more than five years practical experience in the preparation of these injections for hospital use.

During the progress of this paper the writer was called upon to supply 50 mil ampoules of 50 per cent. dextrose solutions. In accordance with official directions these were autoclaved. On removal from the autoclave the solutions were deep yellow in color due to caramelization, and would obviously have been condemned by the surgeons for whose use they were intended. The Tyndallization process in conjunction with aseptic methods of preparation was therefore applied for all concentrated solutions of dextrose. A series of solutions of 5, 10, 15, 20, 25, 30 and 50 per cent. strength was prepared and sterilized in the autoclave. Above a concentration of 10 per cent. the solutions were colored, the intensity increasing with concentration. The color is also influenced by the reaction of the glass, and if this is alkaline the degree of caramelization is considerably increased. It would therefore seem advisable for the British Pharmacopœia to stipulate a limit of 10 per cent. concentration for solutions to be sterilized in the autoclave. Steaming of a 30 per cent. solution for an hour produces no visible coloration and, as

results have shown, the solution has been sterile even when prepared under conditions of gross contamination. The change in reaction is also greater in the autoclaved solution as shown by the following results.

TABLE IX.—pH OF DEXTROSE SOLUTIONS

Concentration	pH before sterilisation	pH after steaming	pH after autoclaving
5 per cent.	7.0	6.0	4.6
30 per cent.	6.6	4.6	4.4

Suggestions have been made (5) that sufficient di-sodium hydrogen phosphate should be added to buffer solutions of dextrose to maintain a pH of 7.

SUMMARY.

1. Bacteriological experiments have been made on a number of hypodermic injections showing the results obtained when solutions are prepared under normal conditions.

2. It has been shown that, even in the presence of contamination much worse than that obtaining during any dispensing operations, sterile products can readily be obtained by exposure to streaming steam at atmospheric pressure for one hour.

3. The instructions prescribed by the British Pharmacopœia for sterilization by Tyndallization are inadequate and sterility is not assured.

4. When solutions are incubated during the intervals between successive heatings, the nature of most of the common solutes is such that spores do not pass into the vegetative state and are consequently not destroyed when exposed to a temperature of 80 degrees C. and maintained at 80 degrees to 85 degrees C. for one hour on three successive days.

5. The pH of a number of common hypodermic solutions has been determined before and after steaming for one hour and indications of decomposition of the solute are evidenced in certain cases.

6. The sterilization of solutions of dextrose has been investigated and results suggest that concentrations in excess of 10 per cent. should not be sterilized by autoclaving at 10 pounds per square inch pressure.

My thanks are due to Professor C. C. Okell, Professor of Bacteriology in the University College Hospital Medical School,

under whose supervision the work has been done; to Dr. C. H. Hampshire for the information concerning the sterilization requirements of other national pharmacopœias; and to Mr. H. Berry, who has provided facilities for checking many of the above results in the dispensing classes under his control.

REFERENCES

1. *The Theory of Disinfection, A System of Bacteriology, Med. Res. Council, Vol. I.*
2. *The Extra Pharmacopœia, 20th Ed., Vol. I.*
3. Todd and Sillar, *Pharm. J.*, 1934, 132, 333.
4. Pfister, *Pharm. Ztg. Berl.*, 1933, 78, 1228.
5. Cooper and Haldane, *Med. J. Austral.*, 1932, 1, 736.

SCIENTIFIC AND TECHNICAL ABSTRACTS

Compiled by Arthur Osol, Ph. D.

Ergoclavine, A New Specific Ergot Alkaloid. W. Küssner. *Arch. Pharm. Berl.* 272, 503 (1934). Through *Quart. Journ. Pharm. Pharmacol.* 7, 618 (1934). The author has obtained from ergot a new alkaloid, ergoclavine, crystallizing in long rectangular plates. The anhydrous base, which is very hygroscopic, melts at 177 degrees to 178 degrees C., and has $[\alpha]_D^{20} = +124^\circ$ (1 per cent. in chloroform). Though these figures are close to those for sensibamine, ergoclavine is distinguished from the latter by the fact that it may be recrystallized repeatedly from alcohol or acetone without decomposition. Elementary analysis gave results corresponding to the formula $C_{31}H_{39}N_5O_6$, though, as with other ergot alkaloids, the difficulty of obtaining the material completely anhydrous is a possible source of error. Titration corresponded to a molecular weight of 573. All samples of ergot examined were found to contain ergoclavine, the yields being 16 to 20 per cent. of the total alkaloids for Spanish ergot; 20 per cent. for Russian; and 6 per cent. for Hungarian. Ergoclavine has, qualitatively and quantitatively, the same specific physiological action as ergotoxine.

Determination of Arsenic in Organic Compounds. F. Monforte. *Ann. Chim. appl. Roma.* 24, 105 (1934). Through *Quart. Journ. Pharm. Pharmacol.* 7, 619 (1934). The arsenic in sodium cacodylate can be readily determined in the following way. Melt about 2 grams of sodium hydroxide in a silver or nickel crucible and to the liquid, kept at a temperature of about 350 degrees C., add gradually about 0.1 gram of the cacodylate. When the reaction has ceased, cool and dissolve the contents of the crucible in hydrochloric acid. Add an excess of Bettendorff's reagent, or 50 grams of Bougault's reagent, keep on a water-bath for an hour, filter off the precipitate of arsenic and wash thoroughly. Transfer the filter paper to a stoppered flask, add 50 cc. of N/10 iodine solution and 2 grams of sodium bicarbonate, leave for an hour, shaking occasion-

ally, until all the arsenic has dissolved, and titrate the excess of iodine with N/10 arsenious oxide. For sodium metharsinite heat about 0.1 gram in a Kjeldahl flask with 5 cc. of fuming nitric acid and evaporate the acid to dryness; repeat this three or four times, then add 15 or 20 cc. of strong hydrochloric acid, evaporate to dryness to remove the last traces of nitric acid, dissolve the residue in hydrochloric acid and continue as for sodium cacodylate.

Efficiency of Sodium Benzoate as a Preservative. O. Weider. *Norsk. farm. Tidsskr.* 42, 154 (1934). Through *Quart. Journ. Pharm. Pharmacol.* 7, 632 (1934). The inefficiency of sodium benzoate as a preservative in neutral or slightly alkaline solutions has already been recorded. The author found that, with an aqueous extract of althea, 0.8 per cent. of sodium benzoate was insufficient to prevent growth of moulds, while 0.02 per cent. of benzoic acid was quite effective for this purpose. Sodium benzoate in a concentration of 0.03 per cent. with the addition of hydrochloric acid, was found to be effective when the pH was reduced to below 4, the actual value depending on a number of factors such as the nature of the solution, etc. At pH 1.5 the extract kept free from moulds without the addition of any preservative.

Determination of Small Quantities of Formaldehyde in Mixtures. Fred Thompson. *Chemist and Druggist*, 121, 523 (1934). Most of the available methods for the determination of this substance only give satisfactory results when the formaldehyde is free. When the formaldehyde and its polymers are mixed with other materials the usual methods are found to be very unreliable. Accordingly, the following method is proposed. Transfer 20 grams of the sample to an 800 cc. Kjeldahl flask, add 25 cc. of water and 10 cc. of dilute sulphuric acid and some paraffin to prevent foaming. Steam distill rapidly (about 110 drops per minute, keeping the volume in the distilling flask fairly constant), catching the first 300 cc. (or 200 cc. if extreme accuracy is not desired) in a volumetric flask. Into each of two 50 cc. volumetric flasks run, accurately, 9 cc. of 0.1 N silver nitrate. Add 6 drops of concentrated

nitric acid. To one of the flasks add an excess of potassium cyanide solution—a 0.1 N solution is most convenient. Make up to the mark with water, shake and filter. Pipette 25 cc. of the filtrate into a flask, add 5 cc. of ferric alum indicator, and titrate the excess of silver nitrate with standard thiocyanate solution. This determination serves as the control.

Pipette 25 cc. of the distillate into a small beaker, add the same amount of cyanide solution that was added to the control, and stir well. Transfer this solution to the other volumetric flask, make up to the mark with water, shake and filter. Pipette 25 cc. of this filtrate into a flask, add 0.5 cc. of ferric alum indicator and titrate the excess of silver nitrate with the standard thiocyanate solution. The percentage of formaldehyde is calculated as follows:

If 300 cc. have been distilled,

$$\frac{(\text{cc. KCNS for sample} - \text{cc. KCNS for control}) \times \text{normality of KCNS} \times 3.6}{100}$$

If 200 cc. have been distilled,

$$\frac{(\text{cc. KCNS for sample} - \text{cc. KCNS for control}) \times \text{normality of KCNS} \times 2.6}{100}$$

The method depends upon the fact that formaldehyde forms an addition product with cyanides and that the cyanide consumed in this reaction does not react with silver nitrate.

Micro-Determination of Acetyl. A. J. Bailey and R. J. Robinson. *Mikrochemie.* 15, 233 (1934). After trying various unsatisfactory modifications, the following procedure was found to yield accurate results. A 5 to 10 mgm. sample is weighed into a small porcelain boat and introduced into a 20 cc. Pyrex flask. 5 cc. of N/25 aqueous sodium hydroxide solution are added to the flask, a micro-reflux condenser attached, and the solution boiled gently for 12 to 35 hours. The time required for complete hydrolysis depends largely upon the character of the acetyl linkage. It has been found that the acetyl-N linkage possesses greater stability than the acetyl-O. After hydrolysis is complete, any alkali adhering to the condenser and rubber stopper is washed into the flask and the excess alkali titrated with N/100 hydrochloric acid using phenolphthalein as the indicator. A blank correction, as determined by a control experi-

ment, is necessary. The authors have found this method to be more accurate than the Pregl-Soltys and Kuhn-Roth methods and report excellent results in the analysis of acetanilid, acetylsalicylic acid, acetyltoluidine, phenacetin, allylisopropylacetylcarbamide and acetan-
iside.

Determination of Acidity in Non-Aqueous Solutions. O. Tomicek and J. Feldman. *Pharm. Journ.* 133, 595 (1934). The determination of acidic substances insoluble in or immiscible with water has hitherto been difficult, because of the uncertain end-points with ordinary indicators, and was impossible with deeply colored solutions. This led to the use of a potentiometric method with hydrogen, quinhydrone, antimony or tellurium electrodes. The tellurium electrode, which has apparently never before been used in determining hydrogen ion concentration, was found to be most satisfactory between the range pH 5.3 to pH 11.5. It proved to be useful in all the instances of non-aqueous acid solutions that were tried. Solutions of benzoic, oxalic, formic and salicylic acids in such solvents as benzene, methyl, ethyl, isopropyl and amyl alcohols, ether, anisole, hexaline, aniline, acetone and turpentine were all used in testing the suitability or limitations of the potentiometric method. The results compared favorably with those obtained using phenolphthalein as indicator. The antimony and tellurium electrodes gave the best results, especially when lithium chloride was added to the N/10 alcoholic potassium hydroxide solution used for titration. Accurate and reproducible results were also obtained with liquid paraffin preparations and with mixed solvents such as are met with in pharmaceutical work. Samples of linseed and poppyseed oils, preferably dissolved in ether, were readily titrated for free acidity and for saponification values. (October Congress of Slav Pharmacists at Belgrade.)

Determination of Iodides and Bromides in Mixtures. R. L. Raigorodskaja and E. S. Binowa. *Pharm. Zentralhalle*, 75, 609 (1934). The authors suggest a method based upon the U. S. S. R. Pharmacopoeia determination of iron iodide in syrup of ferrous iodide. Place 0.1 to 0.2 gram of the mixture in a glass-stoppered flask, add

4 grams of ferric chloride, allow to stand 10 minutes and dilute to 100 cc. with water. Then add 10 cc. of phosphoric acid, mix carefully, add 1 gram of potassium iodide and titrate the liberated iodine with N/10 sodium thiosulphate. Each cc. of this reagent is equivalent to 0.01660 gram of potassium iodide.

The sum of the bromide and iodide is determined by the Volhard method and the bromide content is calculated from the difference between the number of cc. of N/10 silver nitrate used in the former and the number of cc. of N/10 sodium thiosulphate required for the iodide. The results are quite accurate and the method is simple and rapid.

Phenobarbitone and Soluble Phenobarbitone. H. Finnmere. *Australasian Journal of Pharmacy*, 49, 735 (1934). From the number of queries regarding the dispensing of the substances mentioned in the above title, it would appear necessary that pharmacists should come to some agreement among themselves, and with the medical profession, as to the procedure to be adopted when phenobarbitone is ordered in, say, one-half grain doses in mixture form. Accordingly, the following paragraphs have been submitted to the revision committee of the A. P. F. so that order may be brought into the present chaos.

Barbitone and Phenobarbitone: An equal weight of soluble barbitone or soluble phenobarbitone is dispensed if the other ingredients of the prescription are soluble and compatible with them.

Soluble Barbitone and Soluble Phenobarbitone: An equal weight of barbitone or phenobarbitone, respectively, is dispensed if the other ingredients of the prescription comprise substances which will precipitate insoluble barbitone or phenobarbitone. Such are acids, acid salts and acid preparations, e. g., acid syrups and acid infusions (gentian), salts of ammonia and alkaloids. The insoluble barbitone or phenobarbitone will be suspended with compound powder of tragacanth, 5 grains to each fluid ounce.

Syrupus Euphorbiae Compositus. H. Finnmere and E. R. Cole. *Australasian Journal of Pharmacy*, 49, 736 (1934). It has been reported that this syrup gave an unsightly precipitate on standing.

There seemed a possibility that this was caused by the incompatibility of the alkaloids with some tannoid material of the euphorbia, and it was found that this drug does contain tannin, as shown by the green color produced by tincture of ferric chloride. These alkaloidal tannates may be kept in solution by adding a small amount of glycerin. The authors have under observation samples of the syrup containing 5, 10 and 15 minims of glycerin to each fluid drachm, respectively, and will suggest the inclusion of glycerin in the formula if the syrups show no evidence of precipitation on standing.

A Stable Form of Ferrous Chloride. F. E. Carter. *Pharm. Journ.* 7, 389 (1934). The author has performed experiments to show that solutions of ferrous chloride, to which 1 per cent. (W/V) of citric acid has been added, will not oxidize, and will remain bright for a long period. If 5 per cent. of citric acid is used, the solution can be evaporated, and a salt obtained which can be redissolved to give a clear stable solution. Further, ferrous chloride can be made into tablets by using 2 parts of citric acid to 25 parts of ferrous chloride, using dextrose as excipient. These tablets readily dissolve, giving a bright solution.

SOLID EXTRACTS

Hippo-glossus hippoglossus, Linnaeus, is neither a college yell nor the rare name of a rarer orchid. It is piscatorial palaver for the common halibut, whose liver now competes with that of the cod, as a source of vitamin richness. Its expressed oil is tremendously rich in vitamins A and D. Especially A. Think of it—one drop of a good halibut liver oil may contain as much vitamin A as a whole pound of creamery butter and much more vitamin D—which leads us to remark that halibut liver oil is a drug to be cautiously used, and not a common medicine to be indiscriminately used and abused by anyone who wants to give or take it. Even scientific medicine knows too little of these novelties to haphazardly prescribe them.

Some years ago, before the knowledge of vitamins had reached the profession of medicine, a certain proprietary, cod liver oil wine, was alleged to contain in a non-oily and palatable vehicle, some of the virtues of the very oil itself. The then high and mighty authorities in the profession ridiculed and charged as asinine any claim that cod liver oil contained anything else than oil. To be sure, they agreed that cod liver oil expressed from sun-kissed and putrefied livers did contain some alkaloid-like cadaverous compounds, and possibly a little iodine. But nothing else!

How little they knew, as subsequent events have proven.

Yet how strangely that proprietary article continued in sale and used, proving perhaps that the test-tube of time is inevitably firmer in findings than the misguided findings of a science, so-called!

And since liver oils seem to be in good odor at this writing, it may be of interest to note a "brand new use" for oil of the cod. It is now recommended for the treatment of burns. Experimenters have shown that cod liver oil treatment of burns does not influence the primary shock of the burn but is remarkably effective in con-

trolling the secondary infection of large areas. A rapid cleansing of the wound follows its application, and epithelization is stimulated to a degree not seen in any other form of treatment. It is superior to the tannic acid method in that it can be used on the face and in such difficult regions as the buttocks, scrotum and anus. The cod liver oil was used as a salve . . . Attention is called to the fact that commercial cod liver oil is sterile. . . . The experimenters were particularly impressed with the remarkable regeneration of epithelium over large surfaces, such as the entire back.

"Contented cows" have well been called "foster-mothers of the human race." During 1929, for instance, the milk consumption in the United States was 58.7 gallons per capita per year or, on an average, nine pints of milk were consumed by each individual in the United States every week. This is in contrast to 17.6 pounds of butter and 4.6 pounds of cheese per capita per year or 5.4 ounces of butter and 1.4 ounces of cheese per individual per week. Add to this the average per capita consumption during 1929 of condensed milk (2.75 pounds per capita per year), evaporated milk (13.83 pounds per capita per year), and ice cream (twenty-four pints per capita per year), the total will still be a fraction of the per capita consumption of milk as a liquid. A quantity of milk approximately equal to that marketed is estimated to be consumed on farms for feeding humans and animals, etc., and this never enters the market.

For comparison, the consumption of fluid milk per capita per day in the following countries is:

England and Wales—approximately one-half pint; France—one-third pint; Denmark and Sweden—one and one-half pints; and Switzerland, one and four-fifths pints.

One of the latest devices for girth control is the much exploited and tricky Hollywood diet. It consists of a soya bean preparation which is to be used as replacement for one of the big meals of the day. Soya bean meal does contain a little nourishment, but if one really aspires to the shortest distance between the uprights of one's

outline, a more effective, and certainly a cheaper way would be to substitute sawdust for the soya bean. Or better still, do as the Romans did of old with an emetic after dinner, removing the casus belli, yet keeping the tryst with taste.

Nearly one-third of the time, money, and effort expended by the Federal Food and Drug Administration is being devoted to protecting the public from the danger of poisons used in sprays to combat insect pests and diseases that attack fruits and vegetables. This is revealed by W. G. Campbell, chief of the administration in his report to Secretary of Agriculture Wallace which covers the fiscal year ending June 30, 1934.

This agency examined more than 6000 samples of fruits and vegetables to detect poisonous chemicals used in sprays. As a result, fifty-eight consignments containing dangerous quantities of poison residues were seized. Several States were active in making tests to insure safety from poisons. The Director anticipates that it will be necessary to continue the careful watch until science can discover control methods which will make it possible to combat insects and diseases with chemicals which are not dangerous to human beings.

"Crystal crazy" is the charge made by a Federal food authority against a large share of our population. The last year or two has witnessed the development of the new type of medicinal humbug in the marketing of almost innumerable brands of so-called mineral crystals which are essentially laxatives or cathartics and owe their physiological properties to the presence of some well-known therapeutic agent, usually sodium sulphate ("horse-salts"). Medicinal claims of the most extravagant character are made for them and they are sold at outrageous prices. The more adroit manufacturers are careful to restrict such claims to radio and other advertising distributed separately from the interstate package. Where this precaution is taken, action under the Food and Drugs Act is impossible because of lack of jurisdiction.

Probably the earliest book published in the English language on the examination of urine is entitled, "The Judycyall of Uryus," printed in 1512. This is possibly a translation of a manuscript written by Henry Dariel, a monk, in 1379.

"The Differences, Causes and Judgment of Urine," was published in 1541 by Fletcher. He remarked that rhubarb, saffron, cinnamon and broth of cherries, make the urine yellow; cassia, blackish; oil of bay and henbane reddish. He enumerates the sixteen colors of urine, and considers his work complete!

It is hardly believable that nearly fifty million pounds of snuff are manufactured annually in this country although snuff devotees are rarely encountered here.

The use of snuff was once so popular that Pope Urban VIII in 1624 issued an edict of excommunication against its use during church services, because snuffing and sneezing interrupted the religious ceremonies. Its use spread to England, Scotland, and Ireland, and in the eighteenth century snuff taking became the fashion, the lords and ladies taking it in pinches and the scullions and cooks in handfuls. Charles IX, Napoleon, Frederick the Great, Queen Charlotte, Marie Antoinette, all the Georges of England, Lord Nelson, Count von Moltke were devotees of the powdered weed. Dryden, Pope, Swift, Sterne, Addison, Goldsmith, Coleridge, Burns, Doctor Johnson, and other literateurs, took snuff. The method of taking snuff, of opening and tapping the snuff box, became a fine art.

INDEX TO VOLUME 106 OF THE AMERICAN JOURNAL OF PHARMACY

AUTHORS.

	PAGE		PAGE
Amrhein, Florin J.—		Davis, T. Carroll—	
Notes of the Assay of Strychnine Alkaloid in Strychnine Sulphate Tablets	57	Solve This Between Your Jig Saw Puzzles	28
Appel, Theodore B.—		Ender, F.—	
Address at P. C. P. & S. Aerial Pollen Survey Inaugural	389	On the Vitamin Content of Cod-Liver Oil, as Influenced by the Nutritional State of the Cod	102
Bernstein, Mitchell—		Fegley, Homer Burkert—	
The Physician and the Pharmacist of Today	304	The Chemical Composition of Exudates	16
Bethea, Oscar W.—		Fischelis, Robert P.—	
Interdependence of the Various Departments of Science	250	Our Stand on Federal Drug Legislation—Selected Editorial	373
Bird, John C.—		What Organized Pharmacy Is Doing About It	456
Alpha-Dinitrophenol	462	Goehring, Mina F.—	
Calvert, Ralph L.—		The Reaction of Certain Primary Amines With Phenyl Isothiocyanate and O-Tolyl Isothiocyanate	344
Bacteriology and Sanitary Science (Gershenfeld)—Book Review	76	Graham, J. Howard—	
Text Book on Pharmaceutical Chemistry (Bentley and Driver)—Book Review	77	A Contribution to the Photochemistry of Cod Liver Oil..	44
Cook, E. Fullerton—		An Electrical Method for Checking Physiological Salt Solution	325
New Pharmacopoeial Standards for Cod Liver Oil	178	On the Trail of Nicotine During the Smoking Process	256
Davis, H.—		Surface Tension Determinations of Some Barbiturates..	295
The Preparation of Sterile Solutions	474		

	PAGE		PAGE
Griffith, Ivor—		Greene, L. Wilson—	
The Bill Is Dead—Long Live		Schema Zur Mikrochemischen	
the Bill—Editorial	119	Identifikation Von Alkaloiden	
The Blessed Privilege of Wait-		—Book Review	451
ing—Editorial	79	Hanford, W. E.—	
“Book of the Month”—Editorial	1	Studies on Oil of Turpentine:	
The Chemical Formulary—Edi-		Camphor From Pinene-Hy-	
torial	40	drochloride	287
Department Store Hypnotics—		Harding, T. Swann—	
Editorial	323	Curious Remedies—Old and	
Exit Spring Fever—Editorial ..	151	New	211
John Uri Lloyd—Editorial	363	Hartmann, Alexis F.—	
Keeping Up With Science??—		Theory and Practice of Paren-	
Editorial	193	teral Fluid Administration ...	424
Looking Backward—Editorial..	13	Hatcher, Robert A.—	
Modern Drug Encyclopedia and		Cinchonine Solubility	244
Therapeutic Guide—Book Re-		Hepburn, Joseph Samuel—	
view	401	The Chemical Composition of	
The Pharmaceutical Applica-		Exudates	16
tions of Sulfonated Lauryl		The Relative Concentration of	
Alcohol and Kindred Prod-		Esterase and Lipase in Adi-	
ucts	176	pose Tissue	14
Pharmacy and the Committee		Hoch, J. Hampton—	
on Economic Security—Edi-		The Legend and History of	
torial	407	Passiflora	166
Practical Everyday Chemistry		Holm, August—	
—Book Review	403	Utilization of Carbohydrates	
Regulated, Responsible Cos-		and Salts of Organic Acids	
metics—Editorial	283	by C. Diphtheriae in the Pro-	
Regulating the Sale of Medi-		duction of Toxin of High Lf	
cines—Editorial	453	Value	97
Science Marches On—1933—		Horn, David Wilbur—	
Editorial	5	Transition Points of Mixture of	
A Temple of the Future—Edi-		Cow's Butter and Cacao But-	
torial	369	ter	59
Twenty Thousand Doctors		Unsuspected Copper in Domes-	
Can't Be Wrong—Editorial	39	tic Water Supplies	262

PAGE	PAGE
Houghton, R. E.—	Ludlam, Ernest Bowman—
Studies in the Use of 2,4-Di-	Heavy Hydrogen: A Signifi-
nitrophenylhydrazine as a	cant Discovery 72
Quantitative Reagent for Car-	
bonyl Compounds: I. Benzal-	Macht, David I.—
dehyde 62	A Meteorological Adventure in
Kilmer, Fred B.—	Pharmacology 135
The Story of Gardening—Book	
Review 405	Mallory, G. E.—
Kramer, John E.—	A Study of Aloes and a Quan-
Famous Finds by Pharmacists.. 328	titative Determination of
A Few of the Famous Finds by	Some of Its Constituents 81
Pharmacists 134	
The Story of Cerates and Oint-	Moore, Harry McDuffy—
ments 122	The Relative Concentration of
	Esterase and Lipase in Adi-
	pose Tissue 14
LaWall, Charles H.—	
Alchemical Symbols as Used in	Nichols, Adley B.—
Pharmacy 409	Pharmaceutical Formulas, Vol.
British Pharmaceutical Codex—	II—Book Review 404
Book Review 400	
Roman Numerals and Their	Osol, Arthur—
Uses in Early Manuscripts and	Gould's Pocket Medical Dic-
Books 459	tionary—Book Review 450
Wellcome Research Institution,	Organic Chemistry—Book Re-
London, England—Book Re-	view 449
view 450	Scientific and Technical Ab-
	stracts ..67, 110, 140, 181, 227,
	269, 310, 347, 394, 436, 485
Leonard, George F.—	
Utilization of Carbohydrates	Pamcieri, Z.—
and Salts of Organic Acids	Alpha-Dinitrophenol 462
by C. Diphtheriae in the Pro-	
duction of Toxin of High Lf	Perkins, G. W.—
Value 97	Preparation of Tetraiodophthalic
	Anhydride 467
Lingle, Robert M.—	Studies on Oil of Turpentine:
The Assay of Ointment of Boric	Camphor From Pinene-Hy-
Acid 421	drochloride 287
Lloyd, John Uri—	
Before Vitamins—Editorial ... 42	

	PAGE		PAGE
Pipkin, Pauline E.—		Shafer, E. G. E.—	
Impressions of European Pharmacy	171	Alpha-Dinitrophenol	462
Pottebaum, Sister Mary Edward—		Stoneback, William J.—	
The Reaction of Certain Primary Amines With Phenyl Isothiocyanate and O-Tolyl Isothiocyanate	344	Rhus Toxicodendron	374
Poulsion, E.—		Valaer, Peter—	
On the Vitamin Content of Cod-Liver Oil, as Influenced by the Nutritional State of the Cod	102	A Study of Aloes and a Quantitative Determination of Some of Its Constituents	81
Quimba, G. P.—		Viehoever, Arno—	
Preparation of Tetraiodophthalic Anhydride	467	(Chairman) Report of the Committee on Biology of the Pennsylvania Pharmaceutical Association	382
Riesman, David—		Wilson, Margaret Alberta—	
The Past and Future of Pharmacy	264	Transition Points of Mixture of Cow's Butter and Cacao Butter	59
Schwarz, A. John—		Youngken, Heber W.—	
The Standardization of Digitalis	196	A Comparative Study of the Seeds and Spikes of Certain Caulescent Species of Plantago	157

TITLE AND SUBJECT INDEX FOR 1934.

A—Abstract.
M—Medical and Pharmaceutical Note.
S—Solid Extract.

	PAGE		PAGE
Acacia Solution	429, 432	Alkaloids, Action of Ultra-Violet Light on (A)	113
Acetyl, Microdetermination of (A)	487	Alkaloids, Estimation of, by Displacement Titrations (A)	183
Acetylsalicylic Acid, Decomposition in Aqueous Solution (A) ..	67	Allergic Reaction of Dinitrophenol (A)	152
Acidity in Non-Aqueous Solutions, Determination of (A) ...	488	Aloes	388
Aconite Preparations, Deterioration and Stabilization of (A) ..	192	Aloes, A Study of, and a Quantitative Determination of Some of Its Constituents	81
Adipose Tissue, Relative Concentration of Esterase and Lipase in	14	Alpha-Dinitrophenol, Its Purification With Qualitative and Quantitative Data	462
Adrenaline, Solutions of, in Oil (A)	269	Aluminum Ingestion (S)	187
Adsorption Indicators in Argentometric Assays, Use of (A) ..	181	Ambergris (S)	186
Albuminuria, Castellani's Test for (A)	249	American Pharmaceutical Association Officers-Elect for 1934-35	43
Alchemical Symbols as Used in Pharmacy	409	Amidopyrine and Cinchophen Dangerous to Health and Life (M)	444
Alcohol, Determination of, in Anesthetic Chloroform (A)	70	Amines, Primary, Reaction With Phenyl Isothiocyanate and O-Tolyl Isothiocyanate	344
Alcohol Effects (M)	148	Amines, Quinone a Reagent for (A)	436
Alcohol, Ethyl, Medicinal (A) ...	353	Ammonium Chloride, Argentometric Assay of (A)	181
Alcohol vs. Coffee (S)	321	Amygdalin	388
Aldehydes, Other Than Citronellal, Determination of (A)	111	Anabasin—a Powerful Plant Insecticide (M)	149
Alkali Bromides, Are They Therapeutically Equivalent? (A) ...	227	Anstie's Rule (S)	448
Alkali Salicylates and Benzoates, Assay of (A)	228		

PAGE	PAGE
Anthraquinones, Benzyl Alcohol, Method of Analysis 90	Avocado Oil, Its Physical and Chemical Characteristics 75
Antipyretics, Qualitative Separation of (A) 271	Azochloramide (M) 149
Antipyrin, Assay of (A) 312	Bacillus Aertryke in Decomposition of Citric Acid (A) 440
Antipyrine Method for the Evaluation of Medicinal Charcoal (A) 273	Bacteria in Books (S) 322
Antiseptic Action of Ointments and Related Products (A) 188	Bacterial Toxins, Influence of Emulsions on (A) 441
Antiseptic Orvules 37	Bacteriology and Sanitary Science —Book Review 76
Are the Alkali Bromides Therapeutically Equivalent? (A) ... 227	Balsams, Detection of Rosins in (A) 190
Argentometric Assay of Ammonium Chloride (A) 181	Barbiturates, Surface Tension Determinations of Some 295
Argentometric Assays, Use of Adsorption Assays in (A) 181	Barbiturate Test, Quantitative (A) 190
Argyremia, or Silver Sickness (M) 445	Barbituric Acids, Identification of, Therapeutically Important (A) 368
Arsenic (S) 398	Barium and Sulphates, Volumetric Method for Determination of (A) 70
Arsenic, Determination of, in Medicinal Preparations (A) .. 110	Before Vitamins—Editorial 42
Arsenic in Organic Compounds, Determination of (A) 485	Benedict's Solution, Use of, in Micro-Detection of Sugar in Urine and Other Solutions (A) 394
Arsenic, Medicinal, Cancer Cases Reported From 461	Benzoates and Alkali Salicylates, Assay of (A) 228
Arsphenamine and Neoarsphenamine, Identification of (A) ... 113	Benzyl Alcohol Method—Analysis of Anthraquinones 90
Artificial "Spinal Fluid"428, 432	Bibulous Weeds (A) 452
Ascorbic Acid (S) 398	The Bill Is Dead—Long Live the Bill—Editorial 119
Assay of Alkali Salicylates and Benzoates (A) 228	Biology and Pharmacognosy Committee Report of Penna. Pharm. Assoc. 382
Assay of Ointment of Boric Acid 421	Black Plague (S) 448
Assay of Strychnine Alkaloid in Strychnine Sulphate Tablets, Notes on 57	Blessed Privilege of Waiting—Editorial 79
Atabrine 317	
At the Prescription Counter 34	

	PAGE		PAGE
Blood	433	Carbonyl Compounds: I. Benzal-	
"Blood-curdling" (S)	33	dehyde, Studies in the Use of	
Blood Medication—By Skin (M) ..	358	2,4-Dinitrophenylhydrazine as a	
"Book of the Month"—Editorial ..	I	Quantitative Reagent for	62
Book Reviews	76, 362, 400, 449	Casein Paint (S)	187
Boric Acid Ointment, Assay of ..	421	Castellani's Test for Albuminaria	
British Pharmaceutical Codex—		(A)	249
Book Review	400	Catadyne Process of Water	
British Pharmaceutical Codex,		Sterilization (A)	182
Formulas Proposed for (M) ..	116	Catalysts, Deuterium Measures,	
Bromide and Chloride, Detection		Efficiency of (A)	184
With Mercurous Nitrate (A) ..	395	Catnip (S)	355
Bromides, Alkali, Are They Ther-		C. Diphtheriae, Utilization of Car-	
apeutically Equivalent? (A) ...	227	bohydrates and Salts of Organic	
Bromides and Iodides in Mix-		Acids by, in the Production of	
tures, Determination of (A) ..	488	Toxin of High Lf Value	97
Burns, Tannic Acid, Treatment		Celastrus Scandens, Chemical Ex-	
of (M)	114	amination of Entire Plant (A) ..	190
Cacao Butter and Cow's Butter,		Centrifuge Microscope (S)	108
Transition Points of Mixtures ..	59	Cerates and Ointments, The	
Calcium Deposits in Blood Ves-		Story of	122
sels (S)	274	Charcoal, Medicinal, Evaluation	
Calcium Gluconate (A)	269	of, by the Antipyrine Method	
Camphor From Pinene-Hydro-		(A)	273
chloride	287	Chemical Composition of Exu-	
Cancer	383	dates	16
Cancer Cases Reported From Me-		Chemical Constitution of Fowler's	
dicinal Arsenic	461	Solution	96
Capital Punishment (S)	109	Chemical Formulary, The—Edi-	
Carbohydrate Indigestion (S) ...	448	torial	40
Carbohydrates, Utilization of, and		Chemotherapy of Experimental	
Salts of Organic Acids by C.		Tuberculosis of Guinea Pigs	
Diphtheriae in the Production of		(A)	111
Toxin of High Lf Value	97	Chinaman's Chance (S)	185
		Chlorate, Volumetric Determina-	
		tion of (A)	141
		Chloride and Bromide, Detection	
		With Mercurous Nitrate (A) ..	395

	PAGE		PAGE
Chlorohydrin Gas (S)	397	Coffee vs. Alcohol (S)	321
Chloropicrin—A New Fumigant (M)	147	Cold Cream	35
Chocolate (S)	65	Colorimetric Emodin Analysis ..	90
Chymo-Trypsin (P)	23	Comparative Antiseptic Action of Ointments and Related Products (A)	188
Cinchonine Solubility	244	Comparative Study of the Seeds and Spikes of Certain Caules- cent Species of Plantago	157
Cinchophen and Amidopyrine Dangerous to Health and Life (M)	444	Comparison of Karaya Gum and Tragacanth (A)	234
Cinchophen and Liver Atrophy (P)	25	Compound Bismuth Powder	37
Citric Acid, Decomposition of, by Bacillus Aertrycke (A)	440	Compound Elixir of Thyme	34
Cocaine	387	Contented Cows (S)	492
Cocaine Hydrochloride, Effect of Heating on Anaesthetic Power of Buffered Solution of (A) ..	231	Contribution to the Photochemis- try of Cod Liver Oil, A	44
Cockroaches (S)	33	Copper, Unsuspected, in Domestic Water Supplies	262
Cod, Influence of Nutritional State of, on the Vitamin Content of Cod-Liver Oil	102	Cosmetics, Regulated and Re- sponsible (E)	283
Cod Liver Oil (S)	491	Cow's Butter and Cacao Butter, Transition Points of Mixtures..	59
Cod Liver Oil—A Contribution to the Photochemistry of	44	Crystal Crazy (S)	493
Cod Liver Oil as Wound Dress- ing (M)	235	Crystallization Controller for En- gine Oils (S)	399
Cod Liver Oil for Burns (S) ...	491	Curie, Mme.	294
Cod Liver Oil, New Pharmaco- pœial Standards for	178	Curious Remedies—Old and New	211
Cod Liver Oil, Pectin as Emulsi- fier (A)	396	Cyclopropane (S)	146
Cod-Liver Oil, Vitamin Content of, as Influenced by the Nutri- tional State of the Cod	102	Cysteine (P)	24
Cod Livers Mixed With Cocoa (S)	108	Dakin's Solution by Electrolysis (M)	357
Coffee (S)	66	Decomposition of Solutions of Soluble Phenobarbitone (A) ..	231
		Delicate Test for Lead (M)	150
		Department Store Hypnotics— Editorial	323

	PAGE		PAGE
Derris Root (S)	397	Dispensing of Papaverine Hydrochloride (A)	143
Detection of Sucrose in Lactose (A)	227	Displacement Titrations in Estimation of Alkaloids (A)	183
Detergents for China and Glassware (M)	361	Dog Ancestry (S)	448
Deterioration and Stabilization of Aconite Preparations (A)	192	Donovan's Solution, Composition and Stability of (A)	312
Determination of Elemental Sulphur (A)	352	Dreft (S)	322
Deuterium Measures Efficiency of Catalysts (A)	184	Duprene (S)	144
Dextrose Solution	426, 432	Economic Security, Pharmacy and the Committee on—Editorial ..	407
Diabetes (S)	232, 233	Effect of Heating on Anaesthetic Power of Buffered Solutions of Cocaine Hydrochloride (A) ...	231
Diallylbarbituric Acid, and Pyramidon, Molecular Compound of (A)	111	Effects of Tobacco Smoking	455
Diamonds (S)	399	Electrical Method for Checking Physiological Salt Solution ...	325
Digitalin, Standardised Preparation of (M)	148	Electrynx (S)	65
Digitalis, Standardization of	196	Elemental Sulphur, A New Method for the Determination of (A)	67
Dilaudid (M)	238	Emodin, Colorimetric Analysis ..	90
Dinitrophenol, Allergic Reaction of (A)	152	Emulsifiers (Saponins) in Practical Pharmacy (A)	142
Di-Nitro-Phenol and Preparations Must Be Labeled (N)	315	Emulsion Ratios	406
Dinitrophenol, Crude, Purification of	462	Emulsions, Influence on Bacterial Toxins (A)	441
Dinitrophenylhydrazine, 2,4, Studies in the Use of, as a Quantitative Reagent for Carbonyl Compounds: I. Benzaldehyde ..	62	England and the English Language (E)	241
Diodrast	284	England, Joseph Winters—In Memoriam	18
Diphenylamine in the Assay of Saccharated Iron Compounds (A)	349	Ergoclavine (A)	485
Diplogen and Diplon (S)	66	Ergot	386
		Esterase, and Lipase, Relative Concentration in Adipose Tissue	14

PAGE	PAGE
Estimation of Alkaloids by Displacement Titrations (A) 183	Food and Drugs Act (S) 144
Estimation of Official Hypophosphites (A) 140	Formaldehyde, Determination of Small Quantities in Mixtures (A) 486
Ether, U. S. P., Stability of (M) 114	Formulas Proposed for the British Pharmaceutical Codex (M) 116
Ethyl Alcohol and Hydrogen Peroxide, Stability of Mixtures of (A) 311	Fowler's Solution, Chemical Composition of 96
Ethylene Dichloride, Toxicity of (A) 156	France Honors Sir Henry Wellcome 146
Ethylene for Gassing Oranges (S) 276	Fungicides 387
Ethylene Gas, Novel Uses (S) .. 145	Gelatin, Detection in Milk Products (A) 314
Ethylene Glycol in Presence of Glycerin, Test for (A) 371	Girth-Control (S) 274
European Pharmacy, Impressions of 171	Gould's Pocket Medical Dictionary—Book Review 450
Evaluation of Medicinal Charcoal by the Antipyrine Method (A) 273	Glands 382
Exit Spring Fever—Editorial ... 151	Glucose and Normal Saline, A Review of the Causes of Reactions Following the Intravenous Injection of (A) 142
Exudates—Chemical Composition of 16	Glycerin, Test for Ethylene Glycol in (A) 371
Falling Hair (S) 144	Glycerophosphate Syrups, Determination of Phosphorus in (A) 348
Famous Finds by Pharmacists 134, 328	Gravimetric and Volumetric Determination of Mercury as Mercurous Iodate (A) 183
Fats and Waxes, Determination of Saponification Number of (A) 438	Hay Fever and Fly Sprays (A) .. 309
Female Plant Secretions (A) ... 423	Heart Health 435
Ferrous Chloride in Stable Form (A) 490	Heavy Hydrogen: A Significant Discovery 72
Finds by Pharmacists 134	Heavy Water (S) 107, 108
Flies as Test Animals (S) 274	Hexamethylenetetramine, Determination of, by Precipitation of the Double Uranyl and Hexamethylenetetramine Sulphate (A) 436
Fluorides 387	
Fluorides in Drinking Water (S) 33	
Fly Sprays and Hay Fever (A) .. 309	
Fog Dispersion (S) 321	

	PAGE		PAGE
Hippo-Glossus Hippoglossus (S)	491	In Memoriam—Philip H. Johnson	22
Home, Sweet Home—But Where Is It? (M)	446	In Memoriam—Walter Valentine Smith	20
Hormones (S)	232	Insecticide, Plant, Iodine as (A)	351
Hormones	383	Insecticides	387
Hydrogen Peroxide and Ethyl Al- cohol, Stability of Mixtures of (A)	311	Inside Hot Water Bag (S)	185
Hydrogen Peroxide, Reaction With Potassium Iodide in De- termination of Chromium (A)	110	Interdependence of the Various Departments of Science	250
Hypersensitivity	384	Interfacial Tension Measurements Controlling the Refining of Mineral Oils (A)	443
Hypnotics, Department Store— Editorial	323	International Method of Assay of Opium (A)	229
Hypodermic Injections (S)	33	Intravenous Injections of Glucose and Normal Saline, A Review of the Causes of Reactions Fol- lowing (A)	142
Hypophosphites, Official, The Estimation of (A)	140	Investing Time (E)	285
Hypophosphite Syrups, Deter- mination of Phosphorus in (A)	348	Iodides and Bromides in Mix- tures, Determination of (A) ..	488
Identification of Arsphenamine and Neoarsphenamine (A)	113	Iodides, Volumetric Determina- tion of (A)	227
Identification of Therapeutically Important Barbituric Acids (A)	368	Iodine, Elemental, as Plant Insec- ticide (A)	351
Impressions of European Phar- macy	171	Iodine, New Method for Deter- mination of (A)	438
Incompatibilities	38	Iodized Eggs (A)	22
Indicator, New—Nitrazine Yel- low (M)	239	Iron Albuminate Solution, Prep- aration of (A)	352
Injectio Ferri et Arseni	35	Iron in Biological Materials, De- termination of (A)	441
Injectio Peptoni	318	Iron in the Blood (S)	354
Injectio Quininae et Aethylis Car- bamatis	318	Irradiation, Ultraviolet, of Milk (A)	228
Injectio Sodii Morrhuae	318	Isotonic Solutions (A)	272
Injectio Thiosinaminae et Sodii Salicylatis	319	Johnson, Philip H.—In Memoriam	22
In Memoriam—Joseph Winters England	18		

PAGE	PAGE
Karaya Gum and Tragacanth, Comparison of (A) 234	Mercaptans (S) 274
Katadyn Process (S) 354	Merck Manual of Therapeutics and Materia Medica—Book Re- view 362
Keeping Properties of Liquid Ex- tract of Ergot, The (A) 4	Mercurial Ointments, New Method of Analysis (A) 310
Keeping Up With Science??— Editorial 193	Mercuric Iodide, New Volumetric Method for Determination (A) 394
Lactate—Ringer's Solution ..426, 431	Mercurous Iodide, New Volumetric Method for Determination (A) 394
Lactobacillus Acidophilus (S) ... 321	Mercuric Nitrate Ointment (Strong), Improved Method for Assay (A) 311
Lactose, Detection of Sucrose in • (A) 227	Mercuric Oxycyanide (A) 313
Lauryl Alcohol, Sulfonated, and Kindred Products, Pharmaceu- tical Applications of 176	Mercurous Iodate, Gravimetric and Volumetric Determination of Mercury as (A) 183
Lead, A Delicate Test for (M) .. 150	Mercurous Nitrate for Detection of Chloride and Bromide (A) 395
Legend and History of Passiflora 166	Mercury as Mercurous Iodate, Gravimetric and Volumetric De- termination of (A) 183
Legislation, Federal Drug, Our Stand on—Editorial 372	Mercury, Determination of, in Pharmaceutical Preparations (A) 269
Liebig (S) 32	Meteorological Adventure in Pharmacology, A 135
Life and Death 435	Methylene Blue and Picric Acid, Volumetric Determination of (A) 381
Lilly Research Laboratories Dedi- cated 369	Metycaine (M) 116
Linimentum Ammoniatum (A) .. 68	Micro-Detection of Sugar in Urine and Other Solutions by Use of Benedict's Solution (A) 394
Lipase, and Esterase, Relative Concentration in Adipose Tis- sue 14	Milk Products, Detection of Gela- tin in (A) 314
Lloyd, John Uri—Editorial 363	Milk, Ultraviolet Irradiation of (A) 228
Looking Backward—Editorial ... 13	
Lotio Alba, Improved Form (A) 442	
Luminal and Veronal in Their So- dium Salts, A Volumetric Method for the Estimation of (A) 140	
Magnesium, Microanalysis With o-Oxyquinoline (A) 68	
Medicinal Ethyl Alcohol (A) 353	
Medicinal Whisky (M) 114	

	PAGE		PAGE
Mineral Oils, Controlling the Re- fining of, by Interfacial Ten- sion Measurements (A)	443	Nitrogen Trichloride for Gassing Oranges (S)	276
Mistura Quininae Salicylatis	319	Nitron as a Precipitant for Ni- trates (A)	69
Modern Drug Encyclopedia and Therapeutic Guide—Book Re- view	401	Normal Saline and Glucose, A Re- view of the Causes of Reactions Following the Intravenous In- jection of (A)	142
Molecular Compound of Pyra- midon and Diallylbarbituric Acid (A)	111	Nose Sense (S)	355
Money, Cleanliness of (S)	234	Notes on the Assay of Strychnine Alkaloid in Strychnine Sulphate Tablets	57
Monkeys as Drug Addicts (S) ..	186	Nutritional State of Cod, In- fluence on Vitamin Content of Cod-Liver Oil	102
Morphine Made From Poppy Plant Waste (A)	351	Odors, Action of Sunlight and Summer Temperatures on (M) ..	359
Mother's Milk (S)	275	Oil of Turpentine Studies: Cam- phor from Pinene-Hydrochlo- ride	287
Mucilage of Tragacanth, Preser- vation of (A)	348	Ointment Bases, Water-Absorbing Power of (A)	442
Neoarsphenamine and Arsphen- amine, Identification of (A) ...	113	Ointment of Boric Acid, Assay ..	421
Neo-iopax (P)	26	Ointment of Mercuric Nitrate (Strong), Improved Method for Assay (A)	311
Neo-Synephrin Hydrochloride (M)	237	Ointments and Cerates, The Story of	122
New and Non-Official Remedies	236	Ointments and Related Products, Comparative Antiseptic Action of (A)	188
New Law and the Dispensing of Drugs in Japan—Abstract	56	Ointments, Mercurial, New Meth- od of Analysis (A)	310
New Pharmacopoeial Standards for Cod Liver Oil	178	Onion Breath (S)	186
Nicotine (S)	66, 108	Opium Derivatives in the Treat- ment of Common Cold (P) ...	26
Nicotine, Determination of (A) ..	438	Opium, International Method of Assay (A)	229
Nicotine During the Smoking Process, On the Trail of	256		
Nitrazine Yellow—A New Indi- cator (M)	239		
Nitrite, Rapid Method for Accu- rate Determination of Minute Quantities of (A)	112		

	PAGE		PAGE
Oranges Gassed (S)	276	Pharmacists, Famous Finds by ..	328
Organic Acids, Utilization of Salts of, and Carbohydrates, by C. Diphtheria in the Produc- tion of Toxin of High Lf Value	97	Pharmacology, A Meteorological Adventure in	135
Organic Chemistry—Book Review	449	Pharmacopœial Standards, New, for Cod Liver Oil	178
Ortal-Sodium (M)	240	Pharmacy, Alchemical Symbols Used in	409
O-Tolyl Isothiocyanate, and Phenyl Isothiocyanate, Reaction With Certain Primary Amines	344	Pharmacy and the Committee on Economic Security—Editorial..	407
Oxygen in Gases, A Rapid Colori- metric Method for the Detection and Determination of Small Quantities of (A)	71	Pharmacy, Past and Future of ..	264
Palinaesthesia (S)	32	Phenazone, Assay of (A)	312
Papaverine Hydrochloride, The Dispensing of (A)	143	Pheno-barbital (S)	447
Paprika (S)	232	Phenobarbitone and Soluble Phenobarbitone (A)	489
Paraphenetolthiourea (S)	185	Phenobarbitone, Soluble, Decom- position of Solutions of (A) ..	231
Paraphenylene Diamine Eyelash Dyes (S)	144	Phenyl Isothiocyanate and O- Tolyl Isothiocyanate, Reaction With Certain Primary Amines	344
Parenteral Fluid Administration, Theory and Practice of	424	Phenyl-Mercury Compounds (A)	195
Passiflora, Legend and History of	166	Philadelphia College of Phar- macy and Science, Aerial Pollen Survey Begun by	389
Past and Future of Pharmacy ...	264	Philadelphia College of Phar- macy and Science, 112th Annual Commencement	277
Pectin as Emulsifier for Cod Liver Oil (A)	396	Philadelphians Honored	80
Peet-Grady Test (S)	274	Phosphate Syrups, Determination of Phosphorus in (A)	347
Perchloric Acid Oxidation of Medicaments, Determination of Sulphur and Phosphorous in (A)	313	Phosphorus and Sulphur in Med- icaments After Oxidation With Perchloric Acid, Determination of (A)	313
Permutizing Water (S)	322	Phosphorus, Determination of, in Phosphate Hypophosphite and Glycerophosphate Syrups (A)..	347
Pharmaceutical Applications of Sulfonated Lauryl Alcohol and Kindred Products	176	Photochemistry of Cod Liver Oil, A Contribution to	44
Pharmaceutical Formulas, Vol. II —Book Review	404		

	PAGE		PAGE
Physician and Pharmacist of To-day	304	Preparation of Iron Albuminate Solution (A)	352
Physiological Salt Solution, Electrical Method for Checking ...	325	Preparation of Sterile Solutions..	474
Picric Acid and Methylene Blue, Volumetric Determination of (A)	381	Preparation of Tetraiodophthalic Anhydride	467
Pilulae Phenolphthaleini Compositae	319	Preservation of Mucilage of Tragacanth (A)	348
Pinene-Hydrochloride, Camphor From	287	Prevention of Rancidity by Light Exclusion (M)	235
Plantago, A Comparative Study of the Seeds and Spikes of Certain Caulescent Species of	157	Procter International Award ...	363
Poison Ivy	374	"Professional Pharmacy" Reprints	117
Poisons in Sprayed Fruits and Vegetables (S)	493	Progress in the Medical Sciences	23
Poke Root	388	Protactinium (S)	398
Pollen Survey (Aerial) Begun by Philadelphia College of Pharmacy and Science	389	Psyllium, Further Studies of (A)	191
Poppy Plant Waste Source of Morphine (A)	351	Pulvis Barii Sulphatis Compositus	319
Potassium Iodide, Reaction With Hydrogen Peroxide in Chromium Determination (A)	110	Pyramidon (S)	186
Potassium, New Spot Test for (A)	373	Pyramidon, and Diallylbarbituric Acid, Molecular Compound of (A)	111
Practical Everyday Chemistry—Book Review	403	Pyrethrum	387
Precipitation of Alkaloids by Tannins and the Use of Antipyrine in the Detection of Tannins (A)	31	Qualitative Separation of Antipyretics (A)	271
Pregl's Solution of Iodine (A) ..	111	Quinine Lotion for Sunburn	34
Preparation and Storage of Solutions of Tribromethyl Alcohol (A)	27	Quinine Sulphate, Stability of, During Storage (A)	189
		Quinine Sulphate, Water of, Crystallization of (A)	189
		Quinone: A Reagent for Amines (A)	436
		Radium Production and Importation (S)	109
		Rancidity, Prevention of, by Light Exclusion (M)	235

	PAGE		PAGE
Reaction of Certain Primary Amines With Phenyl Isothiocyanate and O-Tolyl Isothiocyanate	344	Salt in Heart (S)	447
Reactions Following Intravenous Injections of Glucose and Normal Saline, Review of the Causes of (A)	142	Saponification Number of Fats and Waxes, Determination of (A)	438
Recovery From Overanesthetization (S)	32	Saponins as Emulsifiers in Practical Pharmacy (A)	142
Regulated, Responsible Cosmetics—Editorial	283	Scheele (S)	32
Regulating the Sale of Medicines—Editorial	453	Schema Zur Mikrochemischen Identifikation Von Alkaloiden—Book Review	451
Relative Concentration of Esterase, and Lipase in Adipose Tissue	14	Science Marches On—1933—Editorial	5
Removal of Tattoo Marks (A) ..	350	Scientific and Technical Abstracts	67, 110, 140
Report of Committee on Drug Market of Pennsylvania Pharmaceutical Association	299	Science, the Interdependence of the Various Departments of ..	250
Reprints "The Professional Pharmacy"	117	Sea Water (S)	354
Review of the Causes of Reactions Following Intravenous Injections of Glucose and Normal Saline (A)	142	Silver Picrate, Considerations of (A)	189
Rhus Toxicodendron	374	Silver Sickness, or Argyremia (M)	445
Ringer's Solution	426, 430	Smells (S)	356
Roman Numerals and Their Uses in Early Manuscripts and Books ..	459	Smells Catalogued (S)	33
Rosins in Balsams, Detection of (A)	190	Smith, Walter Valentine—In Memoriam	20
Rotenone	387, 388	Snuff (S)	494
Rotenone (S)	397	Sodium Benzoate, Efficiency as a Preservative (A)	486
Saccharated Iron Compounds, Use of Diphenylamine in Assay of (A)	349	Sodium Bicarbonate and Salicylate, Discoloration of Mixtures Containing (M)	357
		Sodium Bicarbonate Solution ..	428, 431
		Sodium Dinitrophenate	463
		Sodium Lactate Solution	426, 430
		Sodium Monoxide (M)	445
		Sodium Salicylate and Bicarbonate, Discoloration of Mixtures Containing (M)	357

	PAGE		PAGE
Solid Extracts	32, 65, 107	Sulfonated Lauryl Alcohol and Kindred Products, Pharmaceutical Applications of	176
Solve This Between Your Jig Saw Puzzles	28	Sulphur and Phosphorous in Medicaments After Oxidation With Perchloric Acid, Determination of (A)	313
Soya Bean (S)	492	Sulphur, Elemental, Determination of (A)	352
"Spinal Fluid," Artificial	428, 432	Summer Temperatures, Action on Odors (M)	359
Spot Test for Potassium, New (A)	373	Sunlight, Action on Odors (M) ..	359
Spring Fever (S)	145	Sun Tan Filters (M)	360
Stability of U. S. P. Ether (M) ..	115	"Super Caustic" (M)	445
Stabilization and Deterioration of Aconite Preparations (A)	192	Surface Tension Determinations of Some Barbiturates	295
Standardised Preparation of Digitalin (M)	148	Symbols, Alchemical, as Used in Pharmacy	409
Standardization of Digitalis	196	Syrup of Ferrous Iodide, Stabilization of (A)	191
"Stardom's Hollywood Diet" (S) ..	276	Syrup of Tolu (A)	271
Sterile Solutions, Preparation of (A)	310	Syrupus Euphorbiae Compositus (A)	489
Story of Cerates and Ointments, The	122	Tannic Acid Cream (S)	448
Story of Gardening—Book Review	405	Tannic Acid Treatment of Burns (M)	114
Strychnine Alkaloid in Strychnine Sulphate Tablets, Notes on the Assay of	57	Tattoo Marks, Removal of (A) ..	350
Strychnine Sulphate Tablets, Notes on the Assay of Strychnine Alkaloid in	57	Temple of the Future—Editorial ..	369
Studies in the Use of 2,4-Dinitrophenylhydrazine as a Quantitative Reagent for Carbonyl Compounds: I. Benzaldehyde	62	Test for Ethylene Glycol in the Presence of Glycerin (A)	371
Study of Aloes and a Quantitative Determination of Some of Its Constituents	81	Tetraiodophthalic Anhydride, Preparation of	467
Sucrose, Detection of, in Lactose (A)	227	Textbook on Pharmaceutical Chemistry (Bentley and Driver) —Book Review	77
Sugar, Micro-Detection of, in Urine and Other Solutions (A) ..	394	Theory and Practice of Parenteral Fluid Administration	424
		Therapeutic Color Schemes (S) ..	322

PAGE	PAGE
Thymus Extract, Biological Effects of (N) 316	Ultraviolet Light (S) 447
Thymus Gland (S) 232	Ultraviolet Light, Action on Alkaloids (A) 113
Titanium, Volumetric Determination of (A) 437	Unguentum Methylis Salicylatis Compositum (M) 116
Toad-Sweat (S) 107	Unguentum Potassii Polysulphidi (M) 116
Tobacco (S) 320	Unsuspected Copper in Domestic Water Supplies 262
Tobacco Smoking, Effects of 455	Urea (S) 65
Tobacco Tar (S) 65	Urine, Early Books on (S) 494
Tolu, Syrup of (A) 271	Urine, Micro-Detection of Sugar in (A) 394
Tomatoes, Vitamin C in (M) ... 147	Utilization of Carbohydrates and Salts of Organic Acids by C. Diphtheriae in the Production of Toxin of High Lf Value .. 97
Tonsils (S) 275	Value Analogy (S) 355
Toothpaste Ads (S) 144	Veronal and Luminal in Their Sodium Salts, A Volumetric Method for the Estimation of (A) 140
Toxicity of Ethylene Dichloride (A) 156	Vinyl Ether (P) 23
Toxin of High Lf Value, Utilization of Carbohydrates and Salts of Organic Acids by C. Diphtheriae in the Production of ... 97	Vitamin A 103
Toxins 387	Vitamin A (S) 275
Tragacanth and Karaya Gum, Comparison of (A) 234	Vitamin C in Tomatoes (M) ... 147
Trail of Nicotine During the Smoking Process 256	Vitamin Content of Cod Liver Oil as Influenced by the Nutritional State of the Cod 102
Transition Points of Mixtures of Cow's Butter and Cacao Butter 59	Vitamin C, Synthetic (S) 398
Trout Pancreas (S) 320	Vitamin D 103
Tuberculosis, Experimental, of Guinea Pigs, the Chemotherapy of (A) 111	Vitamin D (S) 398
Turnip Juice (S) 107	Vitamins 383
'Twas Ever Thus (M) 446	Vitamose (P) 25
Twenty Thousand Doctors Can't Be Wrong—Editorial 39	Volumetric and Gravimetric Determination of Mercury as Mercurous Iodate (A) 183
Ultraviolet Irradiation of Milk (A) 228	

	PAGE		PAGE
Volumetric Determination of Chlorate (A)	141	Weeds (S)	233
Volumetric Determination of Iodides (A)	227	Wellcome Research Institution, London, England—Book Review	450
Volumetric Method for Estimation of Luminal and Veronal in Their Sodium Salts (A)	140	Wellcome, Sir Henry, Honored by France	146
Volumetric Method, New, for Determination of Mercuric and Mercurous Iodides (A)	394	What Organized Pharmacy Is Doing About It	456
Water-Absorbing Power of Ointment Bases (A)	442	Whisky, Medicinal (M)	114
Water Flowing Uphill (S)	32	White Mustard	388
Water of Crystallization of Quinine Sulphate (A)	189	Words, Their Meaning and Their Use—Editorial	153
Water Sterilization, Catadyne Process of (A)	182	Wound Dressing, Cod Liver Oil as (M)	235
Water Supplies, Unsuspected Copper in	262	Wrapper for Drugs and Foods (S)	399
Waxes and Fats, Determination of Saponification Number of (A)	438	Yeast	388
		Yield of Extractive in Unstandardized Official Extracts, The (A)	12

